




## Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



En82C



Issued November 2, 1911.

## U. S. DEPARTMENT OF AGRICULTURE,

BUREAU OF ENTOMOLOGY—CIRCULAR No. 100.

L. O. HOWARD, Entomologist and Chief of Bureau.

# AN INDEX TO CIRCULARS 1 TO 100 OF THE BUREAU OF ENTOMOLOGY.

BY

ROLLA P. CURRIE AND ANDREW N. CAUDELL.

4844°—Cir. 100—11——1

WASHINGTON : GOVERNMENT PRINTING OFFICE : 1911.

#### BUREAU OF ENTOMOLOGY.

L. O. Howard, Entomologist and Chief of Bureau.
C. L. Marlatt, Entomologist and Acting Chief in Absence of Chief.
R. S. Clifton, Executive Assistant.
W. F. Tastet, Chief Clerk.

F. H. Chittenden, in charge of truck crop and stored product insect investigations.

A. D. Hopkins, in charge of forest insect investigations.

W. D. Hunter, in charge of southern field crop insect investigations.

F. M. Webster, in charge of cereal and forage insect investigations.

A. L. Quaintance, in charge of deciduous fruit insect investigations.

E. F. Phillips, in charge of bee culture.

D. M. Rogers, in charge of preventing spread of moths, field work.

ROLLA P. CURRIE, in charge of editorial work.

MABEL COLCORD, in charge of library.

## United States Department of Agriculture,

#### BUREAU OF ENTOMOLOGY.

L. O. HOWARD, Entomologist and Chief of Bureau.

## AN INDEX TO CIRCULARS 1 TO 100 OF THE BUREAU OF ENTOMOLOGY.

By Rolla P. Currie and Andrew N. Caudell.

The first circular of the present series, or what was formerly known as the second series, of the Bureau of Entomology was issued in 1891, while Circular 99 was issued in 1908. The circulars of the first series, something over 40 in number, were nearly all merely circular letters and had only a temporary value. Very few of these first-series circulars are now known to exist, and even the titles of most of them are now unknown.

While some of the circulars of the present, or second, series were in the nature of emergency circulars, and were therefore soon discontinued or superseded, yet very many others were popular summaries, containing in condensed form all the important data available concerning the particular insects discussed. They have therefore a distinct value to the student and worker in economic entomology as well as to the farmer, forester, horticulturist, and general reader. Many of the insects treated have not yet been made the subjects of bulletins or been considered in other publications of the department, and much information contained in these circulars is therefore not to be found elsewhere.

To make this information more easily available and to prevent it from being overlooked, the present index has been prepared, and it is hoped that it will be of use to all who read or consult the circulars of the bureau.

For convenience the index itself is preceded by a list of circulars 1 to 100, with their revisions. This list may be considered as complete and final, since the policy of issuing revised editions of circulars, i. e., under the original serial number, has been discontinued and no more such will be published.

Where circulars have gone through one or more revised editions the latest edition is the one indexed.

#### LIST OF CIRCULARS 1 TO 100 OF THE BUREAU OF ENTOMOLOGY.

No. 1. Condensed information concerning some of the more important insecticides. pp. 7. May, 1891.

No. 2. The hop plant-louse [Phorodon humuli Schr.] and the remedies to be used

against it. pp. 7, figs. 5, pl. 1. June, 1891.

No. 3. An important enemy to fruit trees. The San Jose scale [Aspidiotus perniciosus Comst.]: Its appearance in the eastern United States; measures to be taken to prevent its spread and to destroy it. By L. O. Howard. pp. 10, figs. 5. April 4, 1893.

No. 4. The army worm (*Leucania unipuncta* Haw.). By L. O. Howard. pp. 5, figs. 3. June 16, 1894.

No. 5. The carpet beetle or "Buffalo moth" (Anthrenus scrophulariæ L.). By L. O. Howard. pp. 4, fig. 1. September 8, 1894.

Same, revised edition. pp. 4, fig. 1. Issued August 10, 1908.

No. 6. The Mexican cotton-boll weevil (Anthonomus grandis Boh.). By L. O. Howard. pp. 5, figs. 3. April 2, 1895.

Same, Spanish edition. pp. 6, figs. 2. April 2, 1895.

No. 7. The pear-tree psylla (*Psylla pyricola* Foerst.). By C. L. Marlatt. pp. 8, figs. 6. May 1, 1895.

No. 8. The imported elm leaf-beetle (Galerucella luteola Müll.; Galeruca xanthomelwna Schrank). By C. L. Marlatt. pp. 4, fig. 1. May 23, 1895.

Same, revised edition. pp. 6, fig. 1. Issued September 22, 1908.

No. 9. Cankerworms [Paleacrita vernata Peck and (Anisopteryx) Alsophila pometaria Harr.]. By D. W. Coquillett. pp. 4, figs. 4. May 24, 1895.

No. 10. The Harlequin cabbage bug, or calico back (Murgantia histrionica Hahn).

By L. O. Howard. pp. 2, fig. 1. May 24, 1895.

No. 11. The rose-chafer [Macrodactylus subspinosus Fab.]. By F. H. Chittenden. pp. 4, fig. 1. May 21, 1895.

Same, revised. pp. 4, fig. 1. Issued July 6, 1909.

No. 12. The Hessian fly (Cecidomyia destructor Say). By C. L. Marlatt. pp. 4. July 26, 1895.

Same, revised edition. pp. 4. April 13, 1900. (Superseded by No. 70.)

No. 13. Mosquitoes and fleas. By L. O. Howard. pp. 4. February 1, 1896.

Same, revised edition. November 15, 1897.

No. 14. (Revision of No. 6.) The Mexican cotton-boll weevil (*Anthonomus grandis* Boh.). By L. O. Howard. pp. 8, figs. 5. February 12, 1896.

Same, Spanish edition. pp. 8, figs. 5. May 18, 1896.

No. 15. General work against insects which defoliate shade trees in cities and towns. By L. O. Howard. pp. 4. March 6, 1896. (In advance from an article entitled "The shade-tree insect problem in the eastern United States," to be published in the Yearbook of the department for 1895.)

No. 16. The larger corn stalk-borer (*Diatrwa saccharalis* Fab.). By L. O. Howard. pp. 3, figs. 3. August 13, 1896.

No. 17. The peach-tree borer (Sannina exitiosa Say). By C. L. Marlatt. pp. 4, fig. 1. September 30, 1896. (Superseded by No. 54.)

No. 18. (Revision of No. 14.) The Mexican cotton-boll weevil (Anthonomus grandis Boh.). By L. O. Howard. pp. 8, figs. 5. February 4, 1897. Same, Spanish edition. pp. 10.

Same, German edition. pp. 11.

No. 19. The clover mite (*Bryobia pratensis* Garman). By C. L. Marlatt. pp. 4, fig. 1. March 5, 1897.

Same, second edition. pp. 4, fig. 1. July 19, 1897.

No. 20. The woolly aphis of the apple (Schizoneura lanigera Hausmann). By C. L. Marlatt. pp. 6, figs. 2. March 15, 1897.

Same, revised edition. pp. 6, figs. 2. June 23, 1908.

No. 21. The strawberry weevil (Anthonomus signatus Say). By F. H. Chittenden. pp. 7, figs. 4. March 18, 1897.

Same, revised edition. pp. 10, figs. 5. Issued September 19, 1908.

No. 22. The periodical cicada in 1897. By E. A. Schwarz. pp. 4. May 1, 1897.

No. 23. The buffalo tree-hopper (*Ceresa bubalus* Fab.). By C. L. Marlatt. pp. 4, figs. 2. May 10, 1897.

No. 24. The two-lined chestnut borer (Agrilus bilineatus Weber). By F. H. Chittenden. pp. 8, fig. 1. July 14, 1897.

Same, revised edition. pp. 7, fig. 1. Issued January 20, 1909.

No. 25. The ox warble (*Hypoderma lineata* Villers). By C. L. Marlatt. pp. 10, figs. 10. July 19, 1897.

No. 26. The pear slug (Eriocampoides limacina Retzius). By C. L. Marlatt. pp. 7, figs. 4. August 28, 1897.

No. 27. The Mexican cotton-boll weevil in 1897. By L. O. Howard. pp. 7. December 31, 1897.

No. 28. The box-elder plant-bug (*Leptocoris trivittatus* Say). By L. O. Howard. pp. 3, fig. 1. January 8, 1898.

No. 29. The fruit-tree bark-beetle (Scolytus rugulosus Ratz.). By F. H. Chittenden. pp. 8, figs. 5. March 25, 1898.

Same, revised edition. March 25, 1903.

No. 30. The periodical cicada in 1898. By E. A. Schwarz. pp. 3. April 1, 1898.

No. 31. The striped cucumber beetle (*Diabrotica vittata* Fab.). By F. H. Chittenden. pp. 7, figs. 2. April 26, 1898.

Same, revised edition. pp. 6, figs. 2. June 9, 1903.

Same, second revision. pp. 8, figs. 2. Issued May 22, 1909.

No. 32. The larger apple-tree borers. By F. H. Chittenden. pp. 12, figs. 3. July 1, 1898.

Same, revised edition. pp. 12, figs. 3. August 30, 1902.

Same, second revise. pp. 11, figs. 3. April 8, 1905.

Same, third revise. pp. 11, figs. 3. Issued December 10, 1907.

No. 33. Remedial work against the Mexican cotton-boll weevil. By L. O. Howard, pp. 6. July 1, 1898. (Supplementary to circulars 18 and 27.)

No. 34. House ants (Monomorium pharaonis et al.). By C. L. Marlatt. pp. 4, figs. 3. July 6, 1898.

Same, revise. pp. 4, figs. 3. Issued June 12, 1907.

No. 35. House flies (*Musca domestica* et al.). By L. O. Howard. pp. 8, figs. 6. July 11, 1898. (Superseded by No. 71.)

No. 36. The true clothes moths (*Tinea pellionella* et al.). By C. L. Marlatt. pp. 8, figs. 3. July 18, 1898.

Same, revise. pp. 8, figs. 3. Issued January 15, 1908.

No. 37. The use of hydrocyanic-acid gas for fumigating greenhouses and cold frames. By Albert F. Woods and P. H. Dorsett. pp. 10, figs. 3. January 10, 1899.

Same, corrected print. (Compare page 8.) pp. 10, figs. 3. January 10, 1899.

Same, revised edition. pp. 10, figs. 3. October 2, 1903.

Same, second revise. pp. 11, figs. 3. Issued April 3, 1908.

No. 38. The squash-vine borer (Melittia satyriniformis Hbn.). By F. H. Chittenden. pp. 6, figs. 2. April 22, 1899.

Same, revised edition. pp. 6, figs. 2. November 19, 1903.

Same, second revise. pp. 6, figs. 2. Issued September 3, 1908.

No. 39. The common squash bug (*Anasa tristis* De G.). By F. H. Chittenden. pp. 5, figs. 3. May 5, 1899.

Same, revised edition. pp. 5, figs. 3. March 23, 1905.

Same, second revise. pp. 5, figs. 3. Issued July 28, 1908.

No. 40. How to distinguish the different mosquitoes of North America. By L. O. Howard (with "Synoptic tables of the North American mosquitoes," by D. W. Coquillett). pp. 8, figs. 3. February 20, 1900.

No. 41. Regulations of foreign governments regarding importation of American plants, trees, and fruits. By L. O. Howard. pp. 4. August 24, 1900.

No. 42. How to control the San Jose scale. By C. L. Marlatt. pp. 6. October 22, 1900.

Same, revised. pp. 6. May 17, 1902.

Same, second edition. May 26, 1902.

Same, third edition. April 6, 1903.

Same, fourth edition. pp. 8. March 10, 1906.

Same, fifth edition. pp. 7. [March 31] 1909.

No. 43. The destructive green pea louse (*Nectarophora destructor Johns.*). By F. H. Chittenden. pp. 8, figs. 3. May 23, 1901.

Same, second edition [entitled "The pea aphis (Macrosiphum pisi Kalt.)"]. pp. 10, figs. 7. Issued February 25, 1909.

No. 44. The periodical cicada in 1902. By W. D. Hunter. pp. 4, figs. 2. March 13, 1902.

No. 45. A new nomenclature for the broads of the periodical cicada. By C. L. Marlatt. pp. 8. May 1, 1902.

No. 46. Hydrocyanic-acid gas against household insects. By L.O. Howard. pp. 6. June 22, 1902.

Same, revised edition. pp. 7. Issued February 20, 1907. Revised by C. L. Marlatt.
No. 47. The bedbug (Cimex lectularius Linn.). By C. L. Marlatt. pp. 8, figs. 3.
June 15, 1902.

Same, revised edition. pp. 8, figs. 3. Issued February 26, 1907.

No. 48. The house centipede (Scutigera forceps Raf.). By C. L. Marlatt. pp. 4, figs. 2. June 10, 1902.

No. 49. The silver fish (*Lepisma saccharina* Linn.). By C. L. Marlatt. pp. 4, figs. 2. June 20, 1902.

No. 50. The white ant (Termes flavipes Koll.). By C. L. Marlatt. pp. 8, figs. 4. June 30, 1902.

Same, revised edition. pp. 8, figs. 4. Issued January 27, 1908.

No. 51. Cockroaches. By C. L. Marlatt. pp. 15, figs. 5. June 23, 1902.

Same, revised. pp. 14, figs. 5. Issued June 23, 1908.

No. 52. The lime, sulphur, and salt wash. By C. L. Marlatt. pp. 8. February 20, 1903.

Same, second edition. March 18, 1904.

No. 53. The yellow-winged locust (Camnula pellucida Scudd.). By C. B. Simpson. pp. 3, fig. 1. May 12, 1903.

No. 54. The peach-tree borer (Sannina exitiosa Say). By C. L. Marlatt. pp. 6, fig. 1. October 19, 1903.

No. 55. Powder-post injury to seasoned wood products. By A. D. Hopkins. pp. 5, fig. 1. November 20, 1903.

No. 56. The most important step in the cultural system of controlling the boll weevil. By W. D. Hunter. pp. 7. October 10, 1904. (Superseded by No. 95.)

No. 57. The greenhouse white fly (*Aleyrodes vaporariorum* Westw.). By A. W. Morrill, pp. 9, fig. 1. February 15, 1905.

No. 58. Report on the gypsy moth and the brown-tail moth, July, 1904. By C. L. Marlatt. pp. 12, pl. 1, map 1. March 21, 1905.

No. 59. The corn root-worms. By F. H. Chittenden. pp. 8, figs. 3. March 23, 1905.

No. 60. The imported cabbage worm (*Pontia rapæ* Linn.). By F. H. Chittenden. pp. 8, figs. 6. April 28, 1905.

No. 61. Black check in western hemlock. By H. E. Burke. pp. 10, figs. 5. May 24, 1905.

No. 62. The cabbage hair-worm. By F. H. Chittenden. pp. 6, fig. 1. May 17, 1905.

Same, revised edition. pp. 6, fig. 1. Issued July 28, 1908.

No. 63. Root-maggots and how to control them. By F. H. Chittenden. pp. 7, figs. 5. June 6, 1905.

Same, second edition. pp. 7, figs. 5. February 12, 1906.

No. 64. The cottony maple scale (*Pulvinaria innumerabilis* Rathvon). By J. G. Sanders. pp. 6, figs. 4. September 7, 1905.

No. 65. The cotton red spider (*Tetranychus gloveri* Bks.). By E. S. G. Titus. pp. 5, figs. 2. October 30, 1905.

No. 66. The joint-worm (*Isosoma tritici* Fitch). By F. M. Webster. pp. 5, figs. 5, October 20, 1905.

Same, revised edition. pp. 7, figs. 6. Issued September 21, 1908.

No. 67. The clover root-borer (*Hylastinus obscurus* Marsham). By F. M. Webster. pp. 5, figs. 4. December 28, 1905.

No. 68. The tobacco thrips and remedies to prevent "white veins" in wrapper tobacco (Euthrips nicotians Hinds). By W. A. Hooker. pp. 5, figs. 2. February 26, 1906.

No. 69. Some insects affecting the production of red clover seed. By F. M. Webster. pp. 9, figs. 8. April 12, 1906.

No. 70. The Hessian fly (Mayetiola [Cecidomyia] destructor Say). By F. M. Webster. pp. 16, figs. 16. April 14, 1906.

No. 71. (Revision of Circular No. 35.) House flies (*Musca domestica* et al.). By L. O. Howard. pp. 9, figs. 9. March 27, 1906.

Same, revised edition. pp. 10, figs. 9. September 21, 1906.

No. 72. Key to the known larvæ of the mosquitoes of the United States. By HarrisonG. Dyar. pp. 6, fig. 1. April 27, 1906.

No. 73. The plum curculio (Conotrachelus nenuphar Herbst). By Fred Johnson and A. A. Girault. pp. 10, figs. 5. April 14, 1906.

No. 74. The periodical cicada in 1906. By C. L. Marlatt. pp. 5, figs. 3. April 16, 1906.

No. 75. Requirements to be complied with by nurserymen or others who make interstate shipments of nursery stock. By A. F. Burgess. pp. 6. July 5, 1906.

Same, revised edition. pp. 7. Issued July 24, 1908. Same, second revision. pp. 9. Issued August 7, 1909.

No. 76. List of publications of the Bureau of Entomology. Compiled by Mabel Colcord. pp. 21. September 24, 1906.

Same, revised to March 1, 1908. pp. 28.

Same, revised to May 1, 1909. pp. 29.

Same, revised to February 1, 1910. pp. 32.

No. 77. Harvest mites, or "chiggers." By F. H. Chittenden. pp. 6, figs. 3. September 29, 1906.

No. 78. The slender seed-corn ground-beetle (Clivina impressifrons Lec.). By F. M. Webster. pp. 5, figs. 2. October 4, 1906.

No. 79. The brood diseases of bees. By E. F. Phillips. pp. 5. October 3, 1906.

No. 80. The melon aphis (Aphis gossypii Glov.). By F. H. Chittenden. pp. 16, figs. 6. November 14, 1906.

No. 81. The aphides affecting the apple. By A. L. Quaintance. pp. 10, figs. 8. Issued March 9, 1907.

No. 82. Pinhole injury to girdled cypress in the South Atlantic and Gulf States. By A. D. Hopkins. pp. 4, fig. 1. Issued January 2, 1907.

No. 83. The locust borer [Cyllene robiniæ Forst.] and methods for its control. By A. D. Hopkins. pp. 8, figs. 4. Issued February 19, 1907.

No. 84. The grasshopper problem and alfalfa culture. By F. M. Webster. pp. 10, figs. 8. Issued April 6, 1907.

No. 85. The spring grain-aphis (*Toxoptera graminum* Rond.). By F. M. Webster. pp. 7, figs. 3. Issued March 29, 1907. (See also No. 93.)

No. 86. The corn leaf-aphis [Aphis maidis Fitch] and corn root-aphis [Aphis maidiradicis Forbes]. By F. M. Webster. pp. 13, figs. 4. Issued May 6, 1907.

No. 87. The Colorado potato beetle (*Leptinotarsa decemlineata* Say). By F. H. Chittenden. pp. 15, figs. 6. Issued June 3, 1907.

No. 88. The terrapin scale (Eulecanium nigrofasciatum Pergande). By J. G. Sanders. pp. 4, figs. 3. Issued May 14, 1907.

No. 89. The periodical cicada in 1907 (*Tibicen septendecim* L., race tredecim Walsh-Riley). By C. L. Marlatt. pp. 4, figs. 3. Issued May 14, 1907.

No. 90. The white-pine weevil [*Pissodes strobi* Peck]. By A. D. Hopkins. pp. 8, figs 6. Issued June 18, 1907.

No. 91. Note on the occurrence of the North American fever tick [(Boophilus) Margaropus annulatus Say] on sheep. By W. D. Hunter. pp. 3. Issued July 3, 1907.

No. 92. Mites and lice on poultry. By Nathan Banks. pp. 8, figs. 6. Issued September 25, 1907.

No. 93. The spring grain-aphis or so-called "green bug" (Toxoptera graminum Rond.). By F. M. Webster. pp. 18, figs. 7. Issued August 22, 1907.

Same, revised. pp. 22, figs. 10. Issued June 23, 1909.

No. 94. The cause of American foul brood [Bacillus larvæ]. By G. F. White. pp. 4. Issued July 29, 1907.

No. 95. (Revision of Circular No. 56.) The most important step in the control of the boll weevil [Anthonomus grandis Boh.]. By W. D. Hunter. pp. 8. Issued October 3, 1907.

Same, revised edition. pp. 8. Issued September 21, 1908.

Same, French edition. pp. 11. Issued January 21, 1909.

No. 96. The catalpa sphinx (*Ceratomia catalpæ* Bdv.). By L. O. Howard and F. H. Chittenden. pp. 7, figs. 2. Issued December 16, 1907.

No. 97. The bagworm (*Thyridopteryx ephemeræformis* Haw.). By L. O. Howard and F. H. Chittenden. pp. 10, figs. 11. Issued February 6, 1908.

No. 98. The apple-tree tent caterpillar (Malacosoma americana Fab.). By A. L. Quaintance. pp. 8, figs. 4. Issued January 28, 1908.

No. 99. The nut weevils. By F. H. Chittenden. pp. 15, figs. 14. Issued March 6, 1908.

No. 100. An index to Circulars 1 to 100 of the Bureau of Entomology. By Rolla P. Currie and Andrew N. Caudell. pp. i-ii, 1-49. Issued October —, 1911.

### INDEX.

[Two numbers joined by a dash, as 1-39, indicate that the subject is treated continuously throughout the pages noted.]

	Circular No.	Page.	Figure
Α.			
Abies grandis. (See Fir, lowland.)			
lasiocarpa. (See Fir, Alpine.)			
lasiocarpa. (See Fir, Alpine.) Ablerus clisiocampæ, parasite of Malacosoma americana	98 80	5 5	
saccharinum, food plant of Eulecanium nigrofasciatum		2	
Achillea millefolium, flowers, food of Anthrenus scrophularix	5, rev.	$\frac{2}{2}$	
Acholla multispinosa, probable enemy of Balaninus proboscideus and B. rectus.		6	
Adalia bipunctata, enemy of aphides		8	7
Macrosiphum pisi	43, ed. 2	5	
Psylla pyricola	37, rev. 2	7 3	
cuneatum, fumigation with hydrocyanic-acid gas		3	
Aëdes atlanticus, distinctive characters of larva	72	4	
distinctive characters of genus, adults fuscus, distinctive characters of adult.	40	4 7	
larva	72	4	
grossbecki, distinctive characters of larva	. 72	6	
infirmatus, distinctive characters of larva	72	5	
quaylei, distinctive characters of larva. sapphirinus, distinctive characters of adult.	40	6 7	
table of North American species, adults	. 40	7	
tormentor, distinctive characters of larva.	72	. 4	
Ageratum, food plant of Aleyrodes vaporariorum	57 78	4	
Agrilus bilineatus, circular	24, rev.	$1-\hat{7}$	1
remedies.	24, rev.	5-7	
Agropyron occidentalis, food plant of Toxoptera graminum repens, food plant of Mayetiola destructor or related	93, rev.	5	
speciestenerum, food plant of Toxoptera graminum	93, rev.	5	
Agrostis, food plant of Mayetiola destructor or related species.	70	7	
possible food plant of Toxoptera graminum	. 85	4	
Alabama, requirements relative to shipment of nursery stock	75, rev. 2	7 6	
(wood) and oil of lemon against corn root-aphis		13	
Aleyrodes citri on citrus plants in greenhouses	57	4	
packardi on strawberry in greenhouses	57 80	$\frac{4}{12}$	
vaporariorum, circular on tomato.	57	1-9	1
on tomato	37, rev. 2	4	
remedies	37, rev. 2 57	5-9	
Alfalfa, food plant of Bruchophagus funebris Camnula pellucida. Diabrotica 12-punctata.	69	8	
Camnula pellucida	84	1 3	
Hylastinus obscurus	59 67	3	
grasshopper enemies, circular	84	1-10	1-8
Alkali wash against round-headed apple-tree borer	32, rev. 3	6	
		6,7	
Allograpta obliqua, enemy of A phis gossypii.  Macrosiphum pisi.	43, ed. 2	6 5	
Almond, lood plant of Bryona pratensis	19, ea. 4	2	
Alocasia macrorhiza variegata, fumigation with hydrocyanic-acid gas.  Alopecurus geniculatus, food plant of Toxoptera graminum	37, rev. 2 93, rev.	3 5	
Alsike. (See Clover, alsike.) Alum ineffective against rose-chafer	11, rev.	3	
Alyssum, sweet, food plant of Pontia rapæ.	60	4	

	Circular No.	Page.	Figure.
Amaranthus, food plant of Aphis gossypii.	80	5	
Amaranthus, food plant of Aphis gossypii.  probable food plant of Diabrotica 12-punctata.	59	4	
Ambrosia beetles in girdled cypress, remediesinjury to girdled cypress	82 82	3-4	
trifida. (See Ragweed.)		1	
Ameloctonus clisiocampæ, parasité of Malacosoma americana	98	5	
ineffective against strawberry weevil	21, rev.	6 7	
to alleviate pain from centipede bites	48	3	
Amorpha, food plant of Diabrotica 12-punctata.	59 61	3 8	
Amphicyrta simplicipes, association with Cheilosia alaskensis	73	6	
A nasa tristis, circular	39, rev. 2	1-5	1-3
remedies.  A natis ocellata = A natis 15-punctata.	39, rev. 2 87	4 10	
15-punctata, enemy of Leptinotarsa decemlineata	/ 87	10	
Andromeda, food plant of Macrodactylus subspinosus	11, rev. 51, rev.	4 4	
Animal matter, dead, food of cockroaches.  Anisopteryx pometaria, general account.	9	3-4	3-4
remedies.	9	3-4	
Anomalon anale, parasite of Malacosoma americanaexile, parasite of Malacosoma americana	98 98	5 5	
Anopedias error, parasite of Dasyneura leguminicola	69	5	
Anopheles albimanus, status.	40	4 6	
annulimanus perhaps = Culex consobrinus	40	4	
barberi, distinctive characters of larva	72	1	
crucians, distinctive characters of adultlarva	. 40	4 1	
synonyms.	40	5	
distinctive characters of genus, adults	40	4	
franciscanus, distinctive characters of larva	72 40	1 5	
maculipennis, distinctive characters of larva	72	1	
probably = Anopheles quadrimaculatus	40	5 4	
status	40	4	
pictus = Anopheles crucians	40	5	
nigripes, status pictus = Anopheles crucians status. punctipennis, adults	40	$\frac{4}{2,4}$	1
Anophetes memans a synonym	40	5	
distinctive characters of adult	40 72	4	
quadrimaculatus, Anopheles maculipennis a probable syn-	12	1	
onym	40	5	
distinctive characters of adultresting position of adults	40 40	4 3	3
table of North American species, adults	40	4	
unrecognized North American species.	40 72	4-5	
Anophelinæ, distinctive characters of larvæ.  Anthomyia ceparum = Pegomya cepetorum.	63, ed. 2	6	
Anthonomus grandis. (See Weevil, Mexican cotton boll.)		1 10	1.5
signatus, circularremedies	21, rev. 21, rev.	1-10 6-10	1-5
Anthrenus lepidus, not known to attack carpets	5, rev.	2	
scrophulariæ, circular	5, rev.	1-4 3-4	1
remedies	5, rev. 2	3	
Ant, little black. (See Monomorium minutum.)			
prown. (See Lasius niger var. americanus.)			
red. (Seè Monomorium pharaonis.) pavement. (See Tetramorium cæspitum.)			
white ( Termes flavipes), circular hydrocyanic-acid gas as remedy.	50, rev.	1-8	1-4
remedies.	46, rev. 50, rev.	7-8	
Ante or disseminators of San Tose scale	3	6	
carbon bisulphid as insecticide. "cows" destruction in control of corn root-aphis.	86	5 8	
destruction in control of corn root-aphis.	86	12	
enemies of Chrysobothris femorata.  feeding on honeydew from aphides.	32, rev. 3 80	11	
house-infesting species, circular.	34, rev.	1-4	1-3
remedies	34, rev.	4 2	
hydrocyanic-acid gas as remedy	46, rev. 34, rev.	4	
Aonidia aurantii, resin wash as insecticide	1	3-4	
citrinus, resin wash as insecticide	26 707	3-4	
A panteles carpatus, parasite of Tinea pellionella Trichophaga tapetzella.	36, rev. 36, rev.	5	
catalpæ. (See Microplitis catalpæ.)			
clisiocampa, parasite of Malacosoma americana	98 96	5 5	
congregatus, parasite of Ceratomia catalpævar. rufocoxalis, parasite of Malacosoma			
americana	98	5	

	Circular No.	Page.	Figure.
A phelinus mali, parasite of Schizoneura lanigera	20, rev.	6	
A phelinus mail, parasite of Schizoneura lanigera nigritus, parasite of Toxoptera graminum. A phenogaster Julva, enemy of Anthonomus signatus	93, rev.	15	
A phenogaster fulva, enemy of Anthonomus signatus	21, rev.	4	
Applies allecting the apple, circular	81	1-10	1-
remedies	81 27 rov 2	9-10	
A phidius fletcheri, parasite of Macrosiphum pisi.  vashingtonensis, parasite of Macrosiphum pisi.  Aphis, apple. (See Aphis mali.)	37, rev. 2 43, ed. 2	5	
washingtonensis, parasite of Macrosiphum nisi.	43, ed. 2	5	
Aphis, apple. (See Aphis mali.)	20,000		1
apple, rosy. (See Aphis malifolix.) apenx. (See Siphocoryne avenx.) black. (See Aphis gossypii.) brassicx, control by natural enemies host of Lysiphlebus tritici.			;
avenæ. (See Siphocoryne avenæ.)			
black. (See Aphis gossypii.)	00	45.40	
orassica, control by natural enemies	80		
cabbage. (See Aphis brassicx.)	93, rev.	17	
	80	5	
citrulli = Aphis gossypii.  corn leaf. (See Aphis maidis.)  root. (See Aphis maidiradicis.)  cotton. (See Aphis ossypii.)  cucumeris = Aphis gossypii.  European grain. (See Siphocoryne avenæ.)  fitchi = Siphocoryne avenæ.	80	5	
corn leaf. (See A phis maidis.)			
root. (See Aphis maidi-radicis.)			
cotton. (See A phis gossypii.)			
cucumeris = Aphis gossypii	80	5	
European grain. (See Siphocoryne avenæ.)	0.1	0	
fitchi = Siphocoryne avenæ	81	3	
forbesi = Aphis gossypii.  gossypii, circular. host of Lysiphlebus tritici. remedies. maidi-radicis, general account. host of Lysiphlebus tritici. remedies.	80	5 1-16	1-
host of Luciablehue tritici	93, rev.	17	1-
remedies	80	8-16	
maidi-radicis, general account	86	1,4-9	3-
host of Lysiphlebus tritici	93, rev.	17	
remedies	86	9-13	
maidis, general account. host of Lysiphlebus tritici. malifolia, general account remedies. mali, general account.	86	1-3	1-
nost of Lysiphleous tritici	93, rev.	17	
munjonæ, general account	81 81	7-9 9-10	
mali general account	81	5-6	1, 3-
remedies	81	9-10	1,0-
	0.1	0 10	
orange. (See Aphis gossypii.)			
pea. (Seè Macrosiphum pisi.)			
melon. (See Aphis gossypii.) orange. (See Macrosiphum pisi.) pea. (See Macrosiphum pisi.) pomi=Aphis mali. setariæ, host of Lysiphlebus tritici sorbi, food plants. perhaps=Aphis malifoliæ. spring grain. (See Toxoptera graminum.) squash. (See Nectarophora cucurbitæ.) woolly. (See Schizoneura lanigera.) apple. (See Schizoneura lanigera.)	81	5	
setariæ, host of Lysiphlebus tritici	93, rev.	17	
soroi, lood plants	81	7	
enring grain (See Terenters graminum)	81	- (	
squash (See Nectarophora cucurhity)			i
woolly. (See Schizoneura lanigera.)			1
apple. (See Schizoneura lanigera.) Aphycus pulvinariæ, parasite of Pulvinaria innumerabilis			
Aphycus pulvināriæ, parasite of Pulvināria innumerabilis	64	4	
Apple aphides, circular	81	1-10	1-
remedies.	81	9-10	
aphis. (See Aphis mali.)			
character of injury by Conotrachelus nenunhar	73	5	
rosy. (See A phis malifolix.) character of injury by Conotrachelus nenuphar crab, food plant of A phis mali	81	5	
Conotrachelus nenuphar	73	4	
Saperda candida	32, rev. 3	3	
cretata	32, rev. 3 32, rev. 3	8	
Schizoneura lanigera	20, rev.	2	
wild, less injured than other apples by Schizoneura lanigera	20, rev.	2	
food plant of Anisopterur nometaria	20, 167.	3-4	
food plant of Anisopteryx pometaria A phis mali	81	5	
malifoliæ	81	7	
sorbi	81	7	
Bryobia pratensis	19, ed. 2	2	
canker-worms	9	1	
Chrysobothris femorata	32, rev. 3	9	
Conotrachelus nenuphar Diabrotica 12-punctata	73 59	4 3	
longicornis	59	7	
vittata	31, rev. 2	2	
Eulecanium nigrofasciatum,	88	9	
Euproctis chrusorrh $\alpha a$	58	11	
Leptocoris trivittatus	. 28	1	
Macrodactylus subspinosus	11, rev.	2	
Malacosoma americana	98	2 1-3	
Paleacrita vernata	9	1-3	
San Jose scale.	64	1	
Saperda candida	32, rev. 3	3	
cretata	32, rev. 3	8	
Schizoneura lanigera	20, rev.	1-4	
Scolytus rugulosus:	29, rev.	2 3	
Siphocoryne avenæ	81		

	Circular No.	Page.	Figure.
Apple, injury by Ceresa bubalus	23	2	
Apple, injury by Ceresa bubalus. louse, woolly. (See Schizoneura lanigera.) Northern Spy variety resistant to Schizoneura lanigeraroot louse (see also Schizoneura lanigera).	20, rev. 2		
insecticides	1	5	
thorn, food plant of Leptinotarsa decemlineata	87	8	
wild balsam. (See Echinocystis lobata.) worm. (See Carpocapsa pomonella.) Apricot, food plant of Conotrachelus nenuphar.	73	4, 5	` <del></del>
Diabrotica 12-punctata	59	3	
not infested by Aspidiotus perniciosus	29, rev. 3	1	
Aradus sp., mistaken for Cimex lectularius	47, rev. 2		
Aralia filicifolia, fumigation with hydrocyanic-acid gas  Arborvitæ, food plant of Bryobia pratensis  Thurdonterus enhemeratormis	19, ed. 2   97	2 3-4	
Thyridopteryx ephemeræformis.  Arctium, food plant of Aphis gossypii.  Areca lutescens, fumigation with hydrocyanic-acid gas.	80	5	
Areca tutescens, lumigation with hydrocyanic-acid gas	37, rev. 2 92	3 4	
Argas miniatus, general account. Arilus cristatus, enemy of Diabrotica 12-punctata. Pontia rapæ.	59 60	5 6	
range extending northward.  Arizona, requirements relative to shipment of nursery stock.	97 75, rev. 2	1-2	
Arkansas, requirements relative to shipment of nursery stock  Army worm. (See Leucania unipuncta.)	75, rev. 2	2	
Arsenate of lead against apple-tree tent caterpillarbagworm,	98 97	9-10	
bagworm. catalpa sphinx Colorado potato beetle imported cabbage worm.	96 87	$^{6}_{14}$	
imported cabbage worm.	60	6	
rose-chafer	73   11, rev.	9 3	
southern corn root-worm striped cucumber beetle	31, rev. 2	5 6-7	
and bordeaux mixture against rose-chaler	11, rev. 21, rev.	8,9	
striped cucumber beetle and fun- gous diseases	01	0.7	
Arsenic in starch paste against silver fish	31, rev. 2   49	6-7 4	
white, against boll weevil. as insecticide	33	3-4 5-7	
against boll weevil	18 17	6-7	
against boll weevil  Arsenical and lime against peach-tree borer  Arsenicals (see also Arsenate of lead, Arsenic, Arsenite of copper, Arsenite of lime, Arsenoid, green, London purple, Paris green, and Scheele's green).			
agamst apple-tree tent caterpmar	98 97	8 9–10	
bagwormboll weevil	33	1-4	
catalpa sphinx Colorado potato beetle imported cabbage worm	. 96	13–14	
imported cabbage wormpear slug	60 26	6	
plum curculio rose-chafer	73	9-10 3	
Southern corn root-worm	11, rev. 59	5	
strawberry weevil	21, rev. 31, rev. 2	8–9 6	
strawberry weevil striped cucumber beetle yellow-winged locust and Bordeaux mixture against apple-tree tent caterpillar.	53 98	3 8	
Southern corn root-worm.	59	5	
milk of lime against apple-tree tent caterpillar harmlessness when properly applied	98	8 7	
Arsenius acid. (See Arsenic, white.) Arsenite of copper against boll weevil.	33	3-4	
lime with soda against Colorado potato beetle	87	14 5-7	
Arsenites as insecticides Arsenoid, green, against plum curculio. Asclepias, food plant of Diabrotica 12-punctata.	73 59 26	9 3 6	
Ashes against pear slug	20, rev.	4	
striped cucumber beetleineffective against strawberry weevil	31, rev. 2 21, rev.	6 7	
ineffective against strawberry weevil.  Ash, mountain, food plant of Chrysobothris femorata.  Saperda candida.	32, rev. 3 32, rev. 3	9	
Scolytus rugulosus	29, rev.	3 2 5	
Asparagus, food plant of Aphis gossypii.  Diabrotica 12-punctata  Aspidiotus perniciosus. (See Scale, San Jose.)  Astar food plant of Alexades supergrigatum	80 59	3	}
Aster, food plant of Aleyrodes vaporariorum	57 59	2 3	
vittata	31, rev. 2	2	

	Circular		
	No.	Page.	Figure.
Atronates collinei porocito of Palvinaria innarmerabilio	64	4	-
Attopates collinsi, parasite of Pulvinaria innumerabilis.  Attagenus piceus, black carpet beetle	36, rev.	4	
destruction by cold storage	36, rev.	8	
trees, and fruits	41	1	
Autographa brassicæ, the cabbage looper	60	6	
Axle grease against ox warble	26	8-9	
В.			
Baccha babista=Baccha clavata	80	6	
clavata, enemy of Aphis gossypii	80 80	6	
cognata, enemy of $A$ $phis$ $gossypii$ . $fuscipennis$ , enemy of $A$ $phis$ $gossypii$ .	80	6	
lugens, enemy of Aphis gossypii  Bacillus alvei.	80 94	6	
$brandenburgiensis$ probably= $Bacillus\ larvx$	94	3	
$burri$ probably= $\overline{B}acillus\ larvx$ $entomotoxicon$ , bacterial disease of $A$ nasa $tristis$	0 94	3 4	
larvæ, cause of American foul brood	39, rev. 2	1-4	
tracheiphilus (see also Neocosmos pora vasinfecta).	1	0	
transmission by Anasa tristis		2	
Bacterial disease of Malacosoma americana	98	6	
wilt disease of eucurbits, cotton, and cowpeas. (See Bacil- lus tracheiphilus and Neocosmospora vasinfecta.)			
Bagging grapes against rose-chafer	11, rev.	. 4	
Bagworm (Thyridopteryx ephemerxformis), circular		1-10 1	
on catalpa	16 15		
remedies	97	8-10	
Bakeries, infestation by cockroaches	51, rev. 99	7 12,14–15	
remedies	. 99	14-15	
obtusus, general accountremedies	99	15 15	
proboscideus, general account	99	2-4, 5-6	2-4, 8, 9 (1,
remedies	99	6,9–12	4,6)
rectus, general account	. 99	2, 4–6	5-7,9 (2,3,5)
remedies	. 99	6,9-12	
Balsam apple, wild. (See Echinocystis lobata.) Banding against fall canker-worm	. 9	4	
peach-tree borer	5 17	3 3–4	
_	11 94	2,3	
spring canker-worm.  Banks, Nathan, circular, "Mites and Lice on Poultry"	. 92	1-8	1-6
Bark beetle fruit. (See Ścolytus rugulosus.) tree. See Scolytus rugulosus.)			
larger fir-tree. (See Hylesinus granulatus.)			
western hemlock. (See Hylesinus n. sp.) maggot, Alpine fir	. 61	9,10	
hemlock. (See Cheilosia alaskensis.) lowland fir. (See Cheilosia hoodianus.)	1	-,	
lowland fir. (See Cheilosia hoodianus.) Sitka spruce	. 61	9,10	
yellow pine, injury to Pinus ponderosa and P. mur-		0,10	
Barley, food plant of Aphis maidis.	. 61	9 2	
(Cecidomyia) Mayetiola destructor		1	
	05	7 4	
Toxoptera graminum	93, rev.	2, 5	
little. (See Hordeum pusillum.)		7	
Barriers against fruit-tree bark-beetle round-headed apple-tree borer.	29, rev. 3 32, rev. 3	6	
Basket worm. (See Bagworm.)			
Bassus latatorius, parasite of syrphus flies that prey upon Macro- siphum pisi	43, ed. 2	5	1
siphum pisi. Bean fly, colloquial name for Pegomya fusciceps.	. 63, ed. 2	1	
food plant of Aleyrodes vaporariorum.  A phis gossypii	57 80	2 5	
Ceresa bubalus	. 23	4	
Diabrotica 12-punctatalongicornis	59 59	3,4	
vittata	. 31, rev. 2	2	
Macrodactylus subspinosus		2	
Pegomya fusciceps  Tetranychus gloveri	. 65	3	
Beans as trap plants against striped cucumber beetle	. 31, rev. 2	5-6 1-8	
Bedbug (Cimex lectularius), circular	46, rev.	1,2	
prey of cockroaches probable prey of Scutigera forceps	. 51, rev.	5 2	

	Circular No.	Page.	Figure.
Beech, food plant of Malacosoma americana	98	. 2	
Pulvinaria innumerabilis. probable food plant of Chrysobothris femorata. Bee-escape treatment against American and European foul broods.	64	2	
Propable 100d plant of Chrysopothris jemorata	32, rev. 3	10	
	36, rev.	3-4 5	
Bees, brood diseases, circular.	79	1-5	
Bees, brood diseases, circular feeding upon honeydew from aphides  Beet, food plant of A phis gossypii.  Diabrotica 12-punctata.  Pegomya fusciceps  Beetles, enemies of canker-worms.  predaceous on house-fly larvæ.	80	1	
Beet, food plant of Aphis gossypii.	80	5	
Penomua fuscicens	63, ed. 2	3 2	
Beetles, enemies of canker-worms.	9	$\frac{2}{2}$	
predaceous on house-fly larvæ	35	. 8	
Begonia, food plant of A phis gossypii.  Belgium, laws relative to importation of American plants, trees,	80	5	
Belgium, laws relative to importation of American plants, trees,	41		
and fruits.  Belladonna, food plant of Leptinotarsa decemlineata.  Benzine against bedbug.  carpet beetle. clothes moths.	87	1 8	
Benzine against bedbug	47, rev.	7	
carpet beetle	5, rev.	3	
clothes moths	36, rev.	6	
clover mite	19, ed. 2	4	
fleas.  Benzoin benzoin, food plant of Eulecanium nigrofasciatum  Bergamot, wild. (See Monarda fistulosa.)  Berries, food plants of Macrodactylus subspinosus.  Bicarbonate of soda. (See Soda.)  Bidens, food plant of Diabrotica 12-punctata.  Birch, white, probable food plant of Chrysobothris femorata.  Birds as disseminators of San Jose scale.	55	4	
Benzoin benzoin, food plant of Eulecanium nigrofasciatum	88	2	
Bergamot, wild. (See Monarda fistulosa.)			
Berries, food plants of Macrodactylus subspinosus.	11, rev.	2	
Bicar Donate of Soda. (See Soda.)	59	3	
Birch, white, probable food plant of Chrusobothris femorata.	32, rev. 3	10	
Birds as disseminators of San Jose scale.	3	. 6	
enemies of canker-worms	9	2	
Diabrotica 12-punctata	59	5	
Diabrotica 12-punctata grasshoppers hosts of Cimer spp	47, rev.	14	
malaria thereof carried by mosquitoes	40	1	
Bisulphid of carbon. (See Carbon bisulphid.)			
Biting habit of Stomoxys calcitrans	{ 35	. 1	
	71, rev. 80	$\frac{1}{5}$	
Black aphis, colloquial name for Aphis gossypii.  beetle, colloquial name for Blatta orientalis	51, rev.	1	
Blackberry, food plant of Anthonomus signatus	21, rev.	2	
Macrodactylus subspinosus	11, rev.	4	
Blackbirds, enemies of grasshoppers.  Black knot. (See Plowrightia morbosa.)	84	4	
Rlacus fuscines, parasite of Scolutus rugulosus	29, rev.	5	
Blankets, horse, food of Trichophaga tapetzella	36, rev.	5	
Blacus fuscipes, parasite of Scolytus rugulosus.  Blacus fuscipes, food of Trichophaga tapetzella. Blarina brevicauda, probable enemy of Isosoma tritici.	66, rev.	6	
Blatta orientalis, general account	51, rev.	3,8-9	. 4
remedies.  Blattella germanica, general account.	51, rev. 51, rev.	3,7,9-10	5
remedies	51, rev.	11-14	
"Blight, American," foreign name for woolly aphis of apple	20, rev.	1	
"Blights," colloquial name for work of chestnut weevil larvæ	99	2 7	
food of Diabrotica longicornis  Blister beetle, striped. (See Epicauta vittata.)	59	- 1	
beetles, the "old-fashioned potato bugs"	87	3	
Blowfly, common name for Calliphora erythrocephala	ſ 35	2	
**	71, rev.	2	
Blueberry. (See Vaccinium sp.) Bluebottle fly. (See Calliphora erythrocephala and Lucilia casar.) small. (See Phormia terranova.) Blue-grass, Kentucky. (See Poa pratensis.)			
Blue-grass, Kentucky. (See Poa pratensis.)	0.0		
Bluejay, enemy of Malacosoma americana	98 54	6	
Boards, partier against peach-tree borer	47, rev.	2	
Bluejay, enemy of Malacosoma americana Boards, barrier against peach-tree borer. Boats, infestation by Cimex Lectularius cockroaches.	51, rev.	3	
Bobwnite, (See Quail.)			
Bæotomus subapterus, parasite of Mayetiola destructor	70	12-13	16
Boll weevil. (See Weevil, Mexican cotton-boll.) Bone, dissolved, ineffective against strawberry weevil.	21, rev	7	
Book bindings and paste eaten by cockroaches	51, rev.	4-5	
lice, hydrocyanic-acid gas as remedy	46, rev.	1	
Books, destruction by Termes flavines	50, rev.	4, 5	
injury by cockroaches	51, rev. 49	4-5 1 2	
Lepisma saccharina Boophilus annulatus on sheep, circular	91	1, 2 1–3	
Borax and sugar against ants	34, rev.	4	
Bordeaux mixture against periodical cicada	89	4	
strawberry weevil	21, rev.	8	
and arsenate of lead against fungous diseases and			
striped cucumber beetle	31, rev. 2	6-7	
	11, rev.	3	
rose-chafer			
	21, rev. 98	8,9	

	Circular No.	Page.	Figure.
Bordeaux mixture and kerosene emulsion against pear-tree psylla		. 8	
Borer, flat-headed apple-tree. (See Enzysobothris femorata.) peach-tree. (See Sannina exitiosa.) round-headed apple-tree. (See Saperda candida.) spotted apple-tree. (See Saperda cretata.)	21, rev.	8,9	
squash-vine. (See Melittia satyriniformis.)  Borers, colloquial name for chestnut weevil larve	99 82	2 1-3	
pin, injury to girdled cypress Bot-fly, colloquial name for Hypoderma lineata. Box elder, food plant of Chrysobothris femorata.  Leptocoris trivittatus.	32, rev. 3 28	1 9 1-3	
Thyridopteryx ephemeræformis	97	2 4	
Brachynus kansanus, enemy of Leptinotarsa decemlineata	21, rev.	10 4	
charus, parasite of Chrysobothris femorata	32, rev. 3	11 5	
pectinator, parasite of Chrysobothris femorata	32, rev. 3	11 7	
poisoned, against grasshoppers.  Bridges, destruction by white ants.  Brimstone (see also Sulphur).		8	
against clover mite.  Brine against powder-post beetles.  British Columbia, laws relative to importation of American plants,	19, ed. 2 55	4 4	
trees, and fruits.  Broccoli, food plant of Murgantia histrionica.	. 41	$\frac{1}{2}$	
Brodætare, Swedish name for Blatta orientalis	51, rev.	9 7	
hordeaceus, food plant of Toxoptera graminumporteri, food plant of Toxoptera graminum	. 93, rev.	4 5	
possible food plant of Toxoptera graminum	85	4 4	
sp., food plant of Toxoptera graminum		. 5	
chilled, overheated, or starved, of bees		5	
"pickle," of bees.  Broom corn, food plant of Aphis maidis. Brown-tail moth. (See Euproctis chrysorrhæa.) Bruchophagus funebris (see also Clover-seed chalcis).	86	5 2	
general accountremedies	. 69	7-9 9	5-8
Brush and cultivator method against pea aphis.  pan method against pea aphis.  Prushing against Colored against pea aphis.	. 43, ed. 2	7-8 8-9 12	
Brushing against Colorado potafo beetle		6	
Bryobia pratensis, circular	. 19, ed. 2	19 1–4	1
remedies  Bud-worm, colloquial name for Diabrotica 12-punctata  tobacco. (See Chloridea virescens and Heliothis obsoleta.)	19, ed. 2 59	4 2	
Buffalo, host of Hypoderma lineata	. 25	1	
Buildings, destruction by white ants.  Bumblebee, agent in pollination of red clover.  Bumelia, food plant of Fulcomium nigrofascintum.	50, rev. 69 88	1,2 1,2 2	
Burdock. (See Arctium.) BURGESS, A. F., circular, "Requirements to be Complied with by Nurserymen or Others who Make Interstate Shipments of Nursery	75 mort 9	1-9	
Stock".  BURKE, H. E., circular, "Black Check in Western Hemlock"	75, rev. 2	1-10 7	1-5
Burlap barriers against fruit-tree bark-beetle	. 98	7-8	
boll weevil	95, rev.	5	
"green bug"		19 3–4	
Hessian flyjoint-worm	70	14 5	
joint-worm powder-post beetles squash bug	. 55	3-4 5	
Burying of nuts in soil against chestnut weevils		11	
rape, colloquial name for Pontia rapæsouthern cabbage. (See Pontia protodice.)	. 60	1	
white, colloquial name for Pontia rapæ	. 60	1	

	Circular No.	Page.	Figure.
С.			
Cabbage aphis. (See Aphis brassicæ.) bug, harlequin. (See Murgantia histrionica.) butterfly. (See Pontia rapæ.)			
southern. (See Pontia protodice.)	. 80	15	
Diabrotica 12-punctata	59 87	3 8	
Murgantia histrionica  Pegomya brassicx	63, ed. 2	$^{1-2}_{3}$	
fusciceps	63. ed 2.1	2 3	
hairworm (See Mermis albicans )	60	4	
looper. (See Autographa brassicæ.) maggot. (See Maggot, cabbage, and Maggot, radish.) rattlesnake, colloquial name for Mermis albicans			
rattlesnake, colloquial name for Mermis albicans  snake, colloquial name for Mermis albicans	62, rev. 62, rev.	1.2	
snake, colloquial name for Mermis albicans worm, colloquial name for Mermis albicans imported. (See Pontia rapæ.) southern. (See Pontia protodice.)	62, rev.	$^{1,2}_{2}$	
"Calico back." (See Murgantia histrionica.) California, requirements relative to shipment of nursery stock	75, rev. 2	. 2	
Calliphora erythrocephala in houses, habits	35 71, rev.	2 2	
Calosoma calidum, enemy of Leptinotarsa decemlineata	87 98	10 6	
Calyptus tibiator. (See Sigalphus tibiator.) Camnula pellucida, circular	53	1–3	1
destructive to grain and grass crops, but little injurious to alfalfa		1	
remedies	53	2-3	
Camphor against ants clothes moths as remedy for "chiggers"	34, rev. 36, rev.	4 6	
Canada, laws relative to importation of American plants, trees, and	77	6	
fruits shipment of nursery stock	75, rev. 2	. 2	
Canker-worm, fall. (See Anisopteryx pometaria.) spring. (See Paleacrita vernata.)			
Canker-worms, circular remedies		2-3.3-4	1-4
Canna, food plant of <i>Diabrotica 12-punctata</i> .  Cannibalism among cockroaches, probability.  larvæ of <i>Pissodes strobi</i> .	59 51, rev.	3	
larvæ of Pissodes strobi	90 50, rev.	7 4	
Cantaloupe (see also Cucurbits, Melons, and Muskmelon). food plant of Diabrotica 12-punctata.	59	3	
loves colleguid pome for Arkingsonneii	31, rev. 2	2 5	
vit  au tata louse, colloquial name for $Aphis$ $gossypii$ .  Cape of Good Hope, laws relative to importation of American plants,	80		
trees, and fruits	41 25	2 9	
periodical cicada, ineffectivestrawberry weevil.	21, rev.	4 8	
as remedy for "chiggers"	77 92	6	
lime against common hen louse oils against ox warble	92 25	- 6	
soap against fruit-tree bark-beetle peach-tree borer	29, rev.	7 3	
round-headed apple-tree borer whitewash against chicken mite	32, rev. 3	6 3	
emulsion against cabbage maggot	63, ed. 2	4	
seed-corn maggotof no value against locust borer	63, ed. 2 83	2 6	
soap, Paris green, lime, and clay against peach-tree borer.	$\left\{\begin{array}{cc} 17 \\ 54 \end{array}\right]$	4	
Carbon bisulphid against ants	34, rev. 63, ed. 2	4 5	2
carpet beetle	5, rev. 99	4 10	
cockroaches imported onion maggot	51, rev. 63, ed. 2	13 7	
little red ant. melon aphis	34, rev.	2 9	
peach-tree borer	54	5	
pecan weevil, possible utility soft-bodied insect pests of cucurbits	99 80	15 16	
subterranean insects	20, rev.	5 4,5	
Cardinal, enemy of Leptinotarsa decemlineata	87		

	Circular No.	Page.	Figure.
Carnations, fumigation with hydrocyanic-acid gas	37, rev. 2	. 4	
black. (See Attagenus piceus.)			
moths. (See Moths, clothes.)	5, rev.	1	
Carpets, food of Anthrenus scrophularix	36, rev.	3	
Tinea pellionella. Trichophaga tapetzella. reported injury by Lepisma saccharina. Carpocapsa pomonella, host of Mermis albicans.	36, rev.	5	
reported injury by Lepisma saccharina	49 (c) row	2 3	
Carpocapsa pomonetta, nost of Mermis atoteans	62, rev. 35	5	
Carrion, breeding place of Musca domestica	171, rev.	3	
Catalpa bignonioides, food plant of Ceratomia catalpæeastern. (See Catalpa bignonioides.) hardy or western. (See Catalpa speciosa.)	96	1	
speciosa, food plant of Ceratomia catalpx	96	1	
sphinx. (See Ceratomia catalpæ.)	00	e	
Catbird, enemy of Ceratomia catalpxgrasshoppers	96 84	6 4	
grasshoppers Caterpillar lime. (See Raupenleim and Dendrolene.)			
Cat. host of Pulex serraticeps	13	3-4	
Catolaccus anthonomi, parasite of Anthonomus signatusincertus, parasite of Anthonomus signatus	21, rev. 21, rev.	4	
Cattle, host of Hypoderma bovis and Hypoderma lineata	25	1-10	
Cattle, host of Hypoderma bovis and Hypoderma tineata.  CAUDELL, ANDREW N., CURRIE, ROLLA P., and, circular, "An Index to Circulars 1 to 100 of the Bureau of Entomology".  Cauliflower, food plant of Diabrotica 12-punctata.  Pegomya brassica:  Pontia rapæ.  Caustia potash and som a susseria de hooded application bores.			
dex to Circulars 1 to 100 of the Bureau of Entomology"	100	i-ii, 1-49	
Pegomua hrassica	63, ed. 2	3	
Pontia rapæ	60	4	
Caustic potash and soap against round-headed apple-tree borer  Ceanothus, food plant of Diabrotica 12-punctata	32, rev. 3 59	6 3	
Cecidomuja destructor (see also Mauetiola destructor).	1	1-4	
circular = Mayetiola destructor remedies = leguminicola. (See Clover-flower midge.)	70	1	
remedies	12, rev.	3-4	
leguminicola. (See Clover-flower midge.) Cedar against clothes moths. Celatoria diabrotice, parasite of Diabrotica 12-punctata	36, rev. 59	6 5	2
vittata	31, rev. 2	4	
Cenocalius populator, parasite of Saperda candida	32, rev. 3	5 1 <del>-4</del> 8	1-2
enemy of house fly	71, rev.	9	9
Ceratomia catalpæ, eircular	48 96	4 1-7	1-2
remedies	96	6-7	
remedies. Cereal products, food of coackroaches. Ceresa bubalus, general account. remedies.	51, rev. 23 23	1-4 4	1-2
Cercis canadensis, food plant of Anthonomus signatus	21, rev.	2	
Cercis canadensis, food plant of Anthonomus signatus	21, rev. 32, rev. 3 32, rev. 3	10	
japonica = Cercis chinensis	32, rev. 3 93, rev.	10 17	
Chalcididæ, parasites of Chrysobothris femorata	32, rev. 3	11	
Chaitophorus Chalcididæ, parasites of <i>Chrysobothris femorata</i> Chalcis fly, enemy of canker-worms	9	2	
ovata, parasite of Thyridopteryx ephemeræformis	97	7 2-8	1
remedies	61	8, 10	1-5
hoodianus, injury to lowland fir. Chenopodium, food plant of Aphis gossypii. Cherry, black Tartarian, not infested by Aspidiotus perniciosus	61	8-9, 10	
Cherry block Texterion not infected by Assistant accommissions	80	5	
food plant of Bryobia pratensis.	3 19	1 2	
Conotrachelus nenuphar Diabrotica 12-punctata	73	4,5	
Diabrotica 12-punctāta	59	3	
vittata Eriocampoides limacina	31, rev. 2 26	1,3	
Eulecanium nigrofasciatum	88	2	
Macrodactylus subspinosus	11, rev.	2	
Sannina exitiosa	{ 17	1	
Scolytus rugulosus	29, rev.	1-2	
ground. (See Physalis.) probable food plant of Chrysobothris femorata wild, food plant of Malacosoma americana Chess, soft. (See Bromus hordeaceus.)		10 2	
Chess, soft. (See Bromus hordeaceus.) Chestnut borer, two-lined. (See Agrilus bilineatus.)			
food plant of Balaninus proboscideus	99	2	
rectus	99	2	
Chrysobothris femorata. weevil, larger. (See Balaninus proboscideus.) lesser. (See Balaninus rectus.)	32, rev. 3	9	
weevils, general account.	99	1-12 6-12	

	Circular No.	Page.	Figure.
Chickadee, enemy of Malacosoma americana	98	6	
Chicken (see also Poultry and Fowls.) enemy of Balaninus caryæ.  Conotrachelus nenuphar mite. (See Dermanyssus gallinæ.)	99 73	$\begin{array}{c} 14 \\ 7 \end{array}$	
tick. (See Argas miniatus.) Chickens, prairie, enemies of grasshoppers Chickwed, food plant of Aphis gossypii. "Chiggers," circular.	84 80	4 5	
colloquial name for harvest mites	77 77	1-6 1	1-3
eradication in the field.  Chigoe. (See Sarcopsylla penetrans.)  Chilocorus himulaerus, enemy of Aphis gossupii.	77 80	6	
Chilocorus bivulnerus, enemy of Aphis gossypii. Leptinotarsa decemlineata. Pulvinaria innumerabilis.	87 64	10	
Chinch bug (Blissus leucopterus), a wheat pest in the United States.  prey of bobwhite or quail  colloquial name for Cimex lectularius	70 87	1 11	
Chinquapin, food plant of Balaninus proboscideus. "Chintze," colloquial name for Cimex lectularius.	47, rev. 99 47, rev.	2 4 2	
Chionasnis sp., on Davallia mooreana	37, rev. 2 29, rev.	2 2 6	4
Chiropachys colon, parasite of Scolytus rugulosus CHITTENDEN, F. H., circular, "Harvest Mites, or Chiggers" "Root Maggots and How to Control	63 od 2	1-6 1-7	1-3
Them". "The Cabbage Hair-worm". "The Colorado Potato Beetle ( <i>Lepti</i> -	63, ed. 2 62, rev.	1-6	1-3
"The Colorado Potato Beetle (Lepti- notarsa decemlineata L.)"" "The Common Squash Bug (Anasa	87	1-15	1-6
"The Common Squash Bug (Anasa tristis De G.)" "The Corn Root-worms" "The Destructive Green Pea Louse	39, rev. 2 59	1-5 1-8	1-3 1-3
( Nectarophora destructor Johns.)'' "The Fruit-tree Bark-beetle (Scolutus	43	1-8	1-3
"The Imported Cabbage Worm (Pon-	29, rev.	1-8	1-5
tia rapz Linn.)". "The Larger Apple-tree Borers" "The Melon Aphis (Aphis gossypii	32, rev. 3	1-8 1-11	1-6 1-3
Glov.)" "The Nut Weevils". "The Pea Aphis ( <i>Macrosiphum pisi</i>	80 99	1-16 1-15	1-6 1-14
Kalt.)'' "The Rose-chafer ( Macrodactylus sub-	43, ed. 2	1-10	1-7
spinosus Fab.)''. "The Squash-vine Borer ( <i>Melittia</i> satyriniformis Hbn.)''.	11, rev. 38, rev. 2	1-14	1 1-2
"The Strawberry Weevil (Anthonomus signatus Say)"	21, rev.	1-10	1-5
"The Striped Cucumber Beetle (Dia- brotica vittata Fab.)"	31, rev. 2	1-8	1-2
"The Two-lined Chestnut Borer (Agrilus bilineatus Weber)" HOWARD, L. O., and, circular, "The Bagwarm	24, rev.	1-7	1
( Thyridopteryx ephemeræfor- mis Haw.) "" "The Catalpa Sphinx (Cera-	97	1-10	1-11
tômia catalpæ Bdv.)" Chloridea virescens, Paris green as remedy	96 68 35	1-7 3-4 6	1-2
Chlorid of lime against house flies.  potash against seed-corn maggot.	71, rev.	6 3	
Chokeberry, food plant of Diabrotica vittata.  Saperda candida  Chrysanthemum, food plant of Alegrodes vaporariorum.	31, rev. 2 32, rev. 3 57	2 3 2	
Diabrotica 12-punctata  Chrysobothris femorata, general account.	59 32, rev. 3	3 8-11	3
remedies	32, rev. 3 80 80	11 6 6	
attenuata, enemy of A phis gossypii larvæ, enemies of Tetranychus gloveri lineaticornis. enemy of A phis gossypii.	65 80	3 6	
nigricornis, enemy of Aphis gossypiioculata, enemy of aphides	80 81	6 8	8
A phis gossypii Macrosiphum pisi Psylla pyricola	43, ed. 2	6 5 5–7	3
plorabunda, enemy of A phis gossypii  Cicada, periodical (Tibicen septendecim), a new nomenclature for broods, circular	80	6	
broods, circularin 1897, circular 1898, circular1898	45 22 30	1-8 1-4 1-3	

	Circular No.	Page.	Figure.
Cicada, periodical (Tibicen septendecim) in 1902, circular	44	1-4	1-2
1906, circular	74	1-5	1-3
17-year broad X 1902	89 44	1-4 1-3	1-3
1906, chediar 1907, chediar 17-year brood X, 1902 XI, 1903 XIV, 1906 XV (Riley),	44	2	
XIV, 1906	74	1-4	
XV (Riley), 1880-1897-			
1914	22	2-3	
XVII (Ri-			
ley), 1881-	20	0.0	
1898 13-year brood VI (Riley),	30	2-3	
1884–1897–			
1910	22	3-4	
VII (Riley), 1885–1898	30	1-2	
XVIII	74	4-5	
XIX, 1907	89	2-4	2
septendecim. (See Cicada, periodical.)			
Cichorium intybus, fumigation with hydrocyanic-acid gas	37, rev. 2	3	
Cinquefoil, yellow-flowered. (See Potentilla canadensis.)			
Circulars of the Bureau of Entomology, index to Nos. 1 to 100	100	i-ii, 1-49	
Cissus discolor, fumigation with hydrocyanic-acid gas	37, rev. 2 57	3 4	
Citrus, food plants of Aleyrodes citri  Clay and cow manure against two-lined chestnut borer.	24, rev.	6	
lime, Paris green, soap, and carbolic acid against peach-tree	17	4	
borer		4	
Cleome, food plant of Pontia rapx	29, rev.	4 6	
Clerus ichneumoneus, enemy of Scolytus rugulosus	29, rev.	6	
Clivina impressifrons, circular.	78	6	1-2
remedies	78	5–6	
planicollis, habitat	78	1	
moths. (See Moths, clothes.)			
Clothing, starched, eaten by Lepisma saccharina	49	1,2	
Clover, alsike, food plant of Hylastinus obscurus	67	8	
crimson, food plant of Bruchophagus funebris	69 59	3	
Diabrotica 12-punctata	43, ed. 2	3,4	
flower midge (Dasyneura leguminicola), general account	69	3-7	2-4
remedies work in relation to	69	6-7	
that of clover-seed			
chalcis	69	2	
work in relation to that of Phytonomus			
punctatus	69	1-2	
food plant of Aphis gossypii  Bryobia pratensis	80	5	
Bryobia pratensis	19, ed. 2		
Diabrotica 12-punctatalongicornis	59 59	3 7	
leaf weevil. (See Phytonomus punctatus.)	99	,	
mammoth, food plant of Hylastinus obscurus	67	3	
mite. (See Bryobia pratensis.) red (see also Trifolium pratense).			
food plant of Bruchophagus funebris	69	8	
Dasyneura leguminicola	69		
Hylastinus obscurus	67	3-4	
Macrosiphum pisi. insects affecting production of seed, circular. root-borer. (See Hylastinus obscurus.)	43, ed. 2 69	1-9	1-8
root-borer. (See Hylastinus obscurus.)	09	1-9	1-0
seed chalcis (see also Bruchophagus funebris).			
	69	2 1	
work in relation to that of clover-flower midge	4	1	
seldom or never eaten by Leucania unipuncta			
seldom or never eaten by Leucania unipuncta	92	- 4	
seldom or never eaten by Leucania unipuncta.  Cluster fly. (See Pollenia rudis.)  Cnemidocoptes gallinæ, parasite of chickens.  mulans, cause of "scaly-leg" in chickens.	92 92	· 4 3	
seldom or never eaten by Leucania unipuncta.  Cluster fly. (See Pollenia rudis.)  Cnemidocoptes gallinæ, parasite of chickens.  mulans, cause of "scaly-leg" in chickens.  Coal oil. (See Kerosene.)	92	3	
seldom or never eaten by Leucania unipuncta.  Cluster fly. (See Pollenia rudis.)  Cnemidocoptes gallinæ, parasite of chickens.  mutans, cause of "scaly-leg" in chickens.  Coal oil. (See Kerosene.)  Cobwebs, food of Tineola biselliella.  Coccinella 9-notata, enemy of Aphis gossypii.	92 36 80	3 4 6,7	3, f, g
seldom or never eaten by Leucania unipuncta.  Cluster fly. (See Pollenia rudis.)  Cnemidocoptes gallinæ, parasite of chickens.  mutans, cause of "scaly-leg" in chickens.  Coal oil. (See Kerosene.)  Cobwebs, food of Tineola biselliella.  Coccinella 9-notata, enemy of Aphis gossypii.  Leptinotarsa decemlineata.	92 36 80 87	3 6.7 10	3,f,g
seldom or never eaten by Leucania unipuncta.  Cluster fly. (See Pollenia rudis.)  Cnemidocoptes gallinæ, parasite of chickens.  mutans, cause of "scaly-leg" in chickens.  Coal oil. (See Kerosene.)  Cobwebs, food of Tineola biselliella.  Coccinella 9-notata, enemy of Aphis gossypii.  Leptinotarsa decemlineata.  Macrosiphum pisi.	92   36 80 87 43, ed. 2	3 6.7 10 5	
seldom or never eaten by Leucania unipuncta.  Cluster fly. (See Pollenia rudis.)  Cnemidocoptes gallinæ, parasite of chickens.  mutans, cause of "scaly-leg" in chickens.  Coal oll. (See Kerosene.)  Cobwebs, food of Tineola biselliella.  Coccinella 9-notata, enemy of A phis gossypii.  Leptinotarsa decemlineata.  Macrosiphum pisi.  Schizoneura lanigera.	92 36 80 87 43, ed. 2 20, rev.	3 6.7 10	
seldom or never eaten by Leucania unipuncia Cluster fly. (See Pollenia rudis.) Cnemidocoptes gallinæ, parasite of chickens. mutans, cause of "scaly-leg" in chickens Coal oil. (See Kerosene.) Cobwebs, food of Tineola biselliella. Coccinella 9-notata, enemy of Aphis gossypii. Leptinotarsa decemlineata. Macrosiphum pisi. Schizoneura lanigera. Toxoptera graminum.	92   36 80 87 43, ed. 2	3 6,7 10 5 6	
seldom or never eaten by Leucania unipuncta Cluster fly. (See Pollenia rudis.) Cnemidocoptes gallinæ, parasite of chickens. mutans, cause of "scaly-leg" in chickens. Coal oil. (See Kerosene.) Cobwebs, food of Tineola biselliella. Coccinella 9-notata, enemy of Aphis gossypii. Leptinotarsa decemlineata. Macrosiphum pisi. Schizoneura lanigera. Tozoptera graminum. sanguinea (see also Cycloneda sanguinea). enemy of Leptinotarsa decemlineata.	92 36 80 87 43, ed. 2 20, rev. 85	3 6.7 10 5 6 5	
seldom or never eaten by Leucania unipuncta Cluster fly. (See Pollenia rudis.) Cnemidocoptes gallinæ, parasite of chickens. mutans, cause of "scaly-leg" in chickens. Coal oil. (See Kerosene.) Cobwebs, food of Tineola biselliella. Coccinella 9-notata, enemy of Aphis gossypii. Leptinotarsa decemlineata. Macrosiphum pisi. Schizoneura lanigera. Tozoptera graminum. sanguinea (see also Cycloneda sanguinea). enemy of Leptinotarsa decemlineata.	92 36 80 87 43, ed. 2 20, rev. 85 87 43, ed. 2	3 4 6.7 10 5 6 5	
seldom or never eaten by Leucania unipuncia Cluster fly. (See Pollenia rudis.) Cnemidocoptes gallinæ, parasite of chickens. mutans, cause of "scaly-leg" in chickens Coal oil. (See Kerosene.) Cobwebs, food of Tineola biselliella. Coccinella 9-notata, enemy of Aphis gossypii. Leptinotarsa decemlineata. Macrosiphum pisi. Schizoneura lanigera. Toxoptera graminum.	92 36 80 87 43, ed. 2 20, rev. 85	3 6.7 10 5 6 5	

	Circular No.	Page.	Figure.
Cocklebur, food plant of Euthrips nicotianæ Cockroach, American. (See Periplaneta americana.) Australian. (See Periplaneta australasiæ.) black. (See Blatta orientalis.) German. (See Blattella germanica.) oriental. (See Blatta orientalis.)	68	2	
Cockroaches, circular	51, rev.	1-14	1-5
enemies of bedbug	{ 47, rev.	7	
hydrocyanic-acid gas as remedy	51, rev. 46, rev.	$\frac{5}{1.2}$	
prey of Scutigera forceps	48	2	
remedies. Codling moth ( Carpocapsa pomonella), arsenicals as insecticides quarantine regulations of New Zealand against introduc-	51, rev. 1	11-14	
tion	41	4	
Colcord, Mabel, circular, "List of Publications of the Bureau of	70		
Colcord, Mabel, circular, "List of Publications of the Bureau of Entomology," rev. to Feb. 1, 1910 Cold. (See Temperature and Weather.) storage against black carpet beetle	76 36, rev.	1-32	
clothes moths.	36, rev.	7-8	
chestnut weevils	99 57	11	
Orthezia insignis	37, rev. 2	3	
fumigation with hydrocyanic-acid gas	37, rev. 2	3	
Collodion, remedy for "chiggers"	77	6	
Comys fusca, parasite of Pulvinaria innumerabilis	75, rev. 2 64	2 4	
Connecticut, requirements relative to shipment of nursery stock Conotrachelus nenuphar. (See Curculio, plum.)	75, rev. 2	2	
Contarinia spp., enemies of A phis gossypii	80 69	6 3	
tritici	96	7	
chestnut weevils	99	12	
Colorado potato beetlegrasshoppers	87 84	15 9	
Copper compounds ineffective against rose-chafer	11, rev.	3	
COQUILLETT, D. W., circular, "Canker-worms"	9	1-4	14
quitões'' Corn, food plant of A phis maidi-radicis	40 86	4-7 1, 4-9	
maidis	86	1-3	
Diabrotica 12-punctata	59	2, 3, 4	
longicornis	31, rev. 2	$\frac{7}{2}$	
vittata Diatræa saccharalis [corn stalk-borer]	16	1-2	
Leucania unipuncta	4	1	
Macrodactylus subspinosus	11, rev. 63, ed. 2	2	
Pegomya fusciceps	65	3	
Toxoptera graminum	85	4	
leaf aphis. (See A phis maidis.) meal against imported cabbage worm.	93, rev.	5 8	
and Paris green against tobacco budworms. root aphis. (See Aphis maidi-radicis.) worm, Southern. (See Diabrotica 12-punctata.) Western or Northern. (See Diabrotica longicornis.)	68	3-4	
worms, circular	59	1-8	1-3
remedies	59	5,7-8	
seed eaten by Clivina impressifrons.	78 86	1-5	
treatment before planting to repel corn root-aphisstalk-borer, larger. (See Diatræa saccharalis.)	00	12	
Cornus, food plant of A phis gossypii	80	5	
Corrosive sublimate against bedbug	47, rev.	7	
clothes moths	36, rev. 59	7 3	
Cosmos, food plant of Diabrotica 12-punctata.  Cotton aphis. (See A phis gossypii.)	00	· ·	
Cotton boll weevil. (See weevil, Mexican cotton-boll.)			
food plant of Anthonomus grandis	6,14,18, 27,33,56,		
	95, rev.		
A phis gossypii	80	4	
Diabrolica 12-punctatalongicornis	59 59	3	
Tetranychus gloveri	65	3 7 1-2	
red spider. (See Tetranychus gloveri.)	27	4	
seed, infestation by boll weevil	14	7-8	
	18	7-8	
stalks, early fall destruction against boll weevil	27	6-7	
	33	4-5	
	56 .		

	Circular No.	Page.	Figure.
	14 27	6,8	
Cotton, volunteer or seppa, and the boll weevil	33 56 95, rev.	3 5 7	
Cottonwood (see also Poplar, Carolina).  Aradus sp. under bark of dead tree.  injury by Ceresa bubalus.	47, rev.	4 2	
Oberea schaumii following work of Ceresa bu- balus	23	2	
location of eggs of Bryobia pratensis.  possible host tree of Chrysobothris femorata.  Covering the plants against squash bug.  strawberry weevil.	19, ed. 2 32, rev. 3 39, rev. 2	3 10 5	
voung plants against striped cucumber beetle	31, rev. 2	6 4-5 4	
Cowbird, enemy of grasshoppers. Cow manure and clay against two-lined chestnut borer.  Cowpea, food plant of Tetranychus gloveri.  Crab apple. (See Apple, crab.)  Cratzgus ozyacantha, food plant of Aphis mali.	65	6 3 5	
spp., food plants of Eulecanium nigrofasciatum.  Creosote against fruit-tree bark-beetle.	81 88 29, rev.	7 2 8	
two-lined chestnut borer. white ants. Criddle mixture against grasshoppers.	24, rev. 50, rev.	7 7 9	
Crop rotation against boll weevil	63, ed. 2	5 4	
corn root-aphis. stalk-borer.	86 16 12, rev.	9 3 4	
Hessian fly. joint-worm western corn root-worm.	66, rev. 59	14 7 8	
"Croton bug," colloquial name for Blattella germanica Crotons, fumigation with hydrocyanic-acid gas. Crow, enemy of Leptinotarsa decemlinada	51, rev. 37, rev. 2	1 3	
Malacosoma americana malarial parasite thereof carried by mosquitoes	40	12 6 1	
Cuckoo, enemy of Leptinotarsa decemlineata. black-billed, enemy of Malacosoma americana yellow-billed, enemy of Malacosoma americana.	98	12 6 6	
Cuckoos, enemies of *Ceratomia catalpægrasshoppers  Cucumber (see also Cucurbits).  beetle, striped. (See *Diabrotica vittata.)	96 84	6 4	
twelve-spotted. (See Diabrotica 12-punctata.) food plant of Aleyrodes vaporariorum. Diabrotica 12-punctata.	57 59	1,2 2,3	
vittata.  Melittia sattyriniformis  prickly. (See Echinocystis lobata.) louse, colloquial name for Aphis gossypii.	31, rev. 2	1 2	
Cucurbit insects, remedies	80 80	5 16	
melon, Pumpkin, Squash, and Watermelon). food plants of A phis gossypii. Diabrotica tongicornis.	80 59	4 7	
Culex annulatus Osten Sacken probably = Culex consobrinus	40	6 6	
boscii probably= Culex pungens status.	40 40	7 6	
consobrinus distinctive characters of adult. synonyms.	40 40 40	5, 6 6	
eonterrens=Psorophora ciliatus. cubensis. probably= Culex pungens.	40 72 40	, 7 3 7	
status damnosus= Culex tæniorhynchus	40 40	6 7	
distinctive characters of adults. excitans, Culex annulatus a probable synonym. distinctive characters of adult.	40	5,6	
excrucians distinctive characters of adult fasciatus, distinctive characters of adult	40 40 40	7 6 5	
synonyms. fatigans, distinctive characters of larva. ferruginosus.		7 3	
$probably = A nopheles \ crucians.$ $frater = Culex \ fasciatus.$	40 40	5 7 6	
status. impatiens= Culex consobrinus.	40 40		

	Circular No.	Page.	Figure.
Culex impiger	40	4, 5, 7	
distinctive characters of adult	40	5	
synonyms	40	7	
implacabilis= Culex impiger	40	7	
incidens= Culex impiger status	40 40	6	
inornatus= Culex consobrinus.	40	6	
mexicanus probably = Culex posticatus	40	7	
status	40	6	
molestus=Psorophora ciliatus	40	7	
mosquito= Culex fasciatus	40	7	
musicus= Culex posticatus	40 40	6	
perturbans	40	7	
distinctive characters of adult	40	6	
pinguis= Culex consobrinus	40	6	
pipiens, distinctive characters of larva	72	3	
posticatus, distinctive characters of adult	40	5	
synonyms	40	7	
provocans probably = Culex stimulans	40	6	
status.  punctor= Culex consobrinus.	40 40	6	
pungens.	40	1,5	
distinctive characters of adult.	40	5, 6	
habits	13	1	
synonyms	40	7	
quinquefasciatus	40	4	
probably = Culex impiger.	40	7	
resting position of adult	40	3	3
restuans, distinctive characters of larva. $rubidus$ probably= $Psorophora\ ciliatus$ .	72 40	4 7	
status.	40	6	
salinarius, distinctive characters of larva	72	2,3	1
signifer	40	7	
distinctive characters of adult	40	6	
sollicitans= Culex tæniorhynchus	40	7	
stimulans, Culex provocans probably a synonym	40	7	
distinctive characters	40 40	5 5-6	
tables of North American species, adults	40	7	
tæniorhynchus, adult	40	2,5	2
distinctive characters of adult	40	5	
synonyms	. 40	7	
tarsalis	40	7	
distinctive characters of adult	40	5,6	
territans, distinctive characters of larva.	72 72	3	
probably= Culex pungens	40	7	
status	40	6	
testaceus probably = Culex consobrinus	40	6	
status	40	6	
triseriatus	40	7	
distinctive characters of adult	40	5 6	
unrecognized North American species.  Culicella dyari, distinctive characters of larva.	40 72	2	
Culicinæ, distinctive characters of larvæ	72	ĩ	
Culiseta absobrinus, distinctive characters of larva	72	4	
consobrinus, distinctive characters of larva	72	4	
Cultural methods against army worm	4	4	
	14	6-8	
	18	6-8 6-7	
boll weevil	27 33	4-6	
·	56	1-7	
	95, rev.	1-8	
buffalo tree-hopper	23	4	
cabbage maggot	63, ed. 2	4	
catalpa sphinx	96	6	
chestnut weevils	99	11-13 6-7	11-12
clover flower midgeroot-borer	69 67	4-5	
seed chalcis.	69	9	
Colorado potato beetle	87	13	
corn root aphis	86		
worm, Southern	59	5	
Western	59	8	
stalk-borer	16	3	
cotton red spider	32, rev. 3	11	
Hav-Headed apple-tice built	29, rev.	6-7	
fruit-tree bark-beetle			
fruit-tree bark-beetle	84	7-8	
fruit-tree bark-beetle grasshoppers "green bug"		6	

	Circular No.	Page.	Figure.
Cultural methods against Hessian fly	∫ 12, rev.	3-4	
	10	13-16	
imported cabbage wormjoint-worm		5-7	
locust borer		7	
melon aphis	. 80	15	
pea aphìspecan weevil	. 43, ed. 2	9-10	
pecan weevil	99 74	14-15	
periodical cicada	73	10	
rose-chafer	. 11. rev.	4	
round-headed apple-tree borer	. 32, rev. 3	7	
squash bug	. 39, rev. 2	5 5	
vine borerstrawberry weevil	. 38, rev. 2 21, rev.	5-6	
striped cucumber beetle	. 31, rev. 2	5-6	
tobacco thrips	. 68	2-3	
two-lined chestnut borer	. 24, rev.	6	
yellow-winged locustCurculio, plum ( <i>Conotrachelus nenuphar</i> ), arsenicals as insecticides	. 53	6	
circular	73	1-10	1-
remedies	. 73	6,7-10	4-
Currant, food plant of Chrysobothris femorata	. 32, rev. 3	10	
Pulvinaria innumerabilis	. 64	2	
JURRIE, ROLLA P., and ANDREW N. CAUDELL, circular, "An in-	100	i-ii-1-49	
Curtains, starched, eaten by Lenisma saccharina	. 49	2	
CURRIE, ROLLA P., and ANDREW N. CAUDELL, circular, "An index to Circulars I to 100 of the Bureau of Entomology".  Curtains, starched, eaten by Lepisma saccharina.  Cuspidors, breeding place of Musca domestica.	. 35	5	
Cutting locust at certain periods to destroy locust borer	- 83	4	
out peach-tree borers	17	4	
round-headed apple-tree borers		5-6 5-6	
useless trees against apple-tree tent caterpillar	. 92,167.5	6-7	
Cycloneda sanguinea (see also Coccinella sanguinea).			
enemy of $A phis gossypii$	. 80	6	
Cyllene robiniæ, circular	. 83	1-8	1
remedies. Cymling (see also Curcurbits and Squash).	. 83	4-8	
food plant of Melittia satyriniformis	. 38, rev. 2	3	
Cyperus, food plant of Diabrotica 12-punctata	. 59	4	,
Cypress, girdled, prevention of insect injury	. 82	3-4	
host tree of Platypus compositus	- 82	2-3	
pinhole injury, circular.	. 82	1-4	
Cyrtoneura stabulans (see also Muscina stabulans).			
in houses	. 35	2,6	
D.			
	85	4	
Dactylis glomerata, food plant of Toxoptera graminum	93, rev.	5	
Dactylis glomerata, food plant of Toxoptera graminum  Dactylopius adonidum on grape	93, rev. 2 37, rev. 2	5 4	
Dactylis glomerata, food plant of Toxoptera graminum  Dactylopius adonidum on grape	93, rev. 2 37, rev. 2	5	
Dactylis glomerata, food plant of Toxoptera graminum  Dactylopius adonidum on grape	93, rev. 2 37, rev. 2	5 4 4 3 4	
Dactylis glomerata, food plant of Toxoptera graminum  Dactylopius adonidum on grape	93, rev. 2 37, rev. 2	5 4 4 3 4 5	
Dactylis glomerata, food plant of Toxoptera graminum  Dactylopius adonidum on grape. remedy.  Dahlia, food plant of Diabrotica 12-punctata.  Daisy, ox-eye, food plant of Macrodactylus subspinosus.  Dandelion, food plant of Aphis gosspyni.  Dandelion of Colloquial name for Blattella germanica.	93, rev. 37, rev. 2 37, rev. 2 59 11, rev. 80 51, rev.	5 4 4 3 4 5 2	
Dactylis glomerata, food plant of Toxoptera graminum  Dactylopius adonidum on grape. remedy.  Dahlia, food plant of Diabrotica 12-punctata.  Daisy, ox-eye, food plant of Macrodactylius subspinosus.  Dandelion, food plant of Aphis gossypii.  "Däne," colloquial name for Blattella germanica.  Daremma catalpæ, same as Ceratomia catalpæ.  Dasymeura leguminicola. (See Clover-flower midge.)	93, rev. 37, rev. 2 37, rev. 2 59 11, rev. 80 51, rev. 96	5 4 4 3 4 5	
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Daisy, ox-eye, food plant of Macrodactylus subspinosus. Dandelion, food plant of A phis gossypti. 'Dâne,'' colloquial name for Blattella germanica. Daremma catalpæ, same as Ceratomia catalpæ. Dasymeura leguminicola. (See Clover-flower midge.) Datura, food plant of A phis gossynii.	93, rev. 2 37, rev. 2 37, rev. 2 37, rev. 2 59 11, rev. 80 51, rev. 96	5 4 4 3 4 5 2 1	
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Daisy, ox-eye, food plant of Macrodactylus subspinosus. Dandelion, food plant of A phis gossypti. 'Dâne,'' colloquial name for Blattella germanica. Daremma catalpæ, same as Ceratomia catalpæ. Dasymeura leguminicola. (See Clover-flower midge.) Datura, food plant of A phis gossynii.	93, rev. 2 37, rev. 2 37, rev. 2 37, rev. 2 59 11, rev. 80 51, rev. 96	5 4 4 3 4 5 2 1	
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape.  remedy.  Dahlia, food plant of Diabrotica 12-punctata  Daisy, ox-eye, food plant of Macrodactylus subspinosus.  Dandelion, food plant of A phis gossypti.  Däne," colloquial name for Blattella germanica  Daremma catalpæ, same as Ceratomia catalpæ.  Dasyneura leguminicola. (See Clover-flower midge.)  Datura, food plant of A phis gossymi.	93, rev. 2 37, rev. 2 37, rev. 2 37, rev. 2 59 11, rev. 80 51, rev. 96	5 4 4 3 4 5 2 1 5 8 4	
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punciata. Dandelion, food plant of Aphis gossypti. Däne,'' colloquial name for Blattella germanica. Daremma catalipæ, same as Ceratomia catalipæ. Dasyneura leguminicola. (See Clover-flower midge.) Datura, food plant of Aphis gossypti. Leptinotarsa decemlineata. probable food plant of Diabrotica 12-punciata. Davallia mooreana, food plant of Chionaspis sp.	93, rev. 37, rev. 2 37, rev. 2 59 11, rev. 80 51, rev. 96 80 87 59	5 4 4 3 3 4 5 2 1 5 8 4 4 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Daisy, ox-eye, food plant of Macrodactylus subspinosus. Dandelion, food plant of A phis gossypii. 'Dâne,'' colloquial name for Blattella germanica. Daremma catalpæ, same as Ceratomia catalpæ. Dasymeura leguminicola. (See Clover-flower midge.) Datura, food plant of A phis gossypii. Leptinotarsa decemtineata. probable food plant of Diabrotica 12-punctata. Davallia mooreana, food plant of Chionospis sp. Deimocerites cancer, distinctive characters of larva Delaware, requirements relative to shipment of nursery stock.	93, rev. 2 37, rev. 2 37, rev. 2 37, rev. 2 59 11, rev. 80 51, rev. 96 87 87 87, rev. 2 75, rev. 2 75, rev. 2	5 4 4 4 4 3 3 4 4 5 5 2 1 1 5 8 8 4 2 2 3 3 2 2	
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Daisy, ox-eye, food plant of Macrodactylus subspinosus. Dandelion, food plant of Aphis gossypii. 'Däne,'' colloquial name for Blattella germanica. Daremma catalpæ, same as Ceratomia catalpæ. Dasyneura leguminicola. (See Clover-flower midge.) Datura, food plant of Aphis gossypii. Leptinotarea decemiineata probable food plant of Diabrotica 12-punctata. Davallia mooreana, food plant of Chionaspis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to pine and spruce.	93, rev. 2 37, rev. 2 37, rev. 2 37, rev. 2 59 11, rev. 80 51, rev. 96 87 87 59 37, rev. 2 75, rev. 2 75, rev. 2 24, rev.	5 4 4 4 3 3 4 4 5 5 2 2 1 1 5 5 8 4 4 2 2 2 2 1 1 2 2 2 1 1 2 2 2 2 2 1	
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Daisy, ox-eye, food plant of Macrodactylus subspinosus. Dandelion, food plant of Aphis gossypii. 'Däne,'' colloquial name for Blattella germanica. Daremma catalpæ, same as Ceratomia catalpæ. Dasyneura leguminicola. (See Clover-flower midge.) Datura, food plant of Aphis gossypii. Leptinotarea decemiineata probable food plant of Diabrotica 12-punctata. Davallia mooreana, food plant of Chionaspis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to pine and spruce.	93, rev. 2 37, rev. 2 37, rev. 2 37, rev. 2 59 11, rev. 80 51, rev. 96 87 87 59 37, rev. 2 75, rev. 2 75, rev. 2 24, rev.	5 4 4 4 3 3 4 5 5 2 2 1 1 5 8 8 4 4 2 2 3 3 2 2 1 9	
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Dandelion, food plant of Macrodactylus subspinosus. Dandelion, food plant of Aphis gossypti. Däne," colloquial name for Blattella germanica. Daremma catalipæ, same as Ceratomia catalipæ. Dasyneura leguminicola. (See Clover-flower midge.) Datura, food plant of Aphis gossypti. Leptinotarsa decemlineata. probable food plant of Diabrotica 12-punctata Davallia mooreana, food plant of Chionaspis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to pine and spruce. sp., injury to Pinus murrayana. ponderosa.	93, rev. 2 37, rev. 2 37, rev. 2 37, rev. 2 11, rev. 80 51, rev. 96 80 87 37, rev. 2 27 75, rev. 2 24, rev. 61	5 4 4 4 3 3 4 4 5 5 2 2 1 1 5 5 8 4 4 2 2 2 2 1 1 2 2 2 1 1 2 2 2 2 2 1	
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Daisy, ox-eye, food plant of Aphis gossypi.  Däne,'' colloquial name for Blattella germanica. Duremma catalpæ, same as Ceratomia catalpæ. Dastura, food plant of Aphis gossypii. Leptinotarsa decemlineata. probable food plant of Diabrotica 12-punctata. Davallia mooreana, food plant of Chionaspis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to pine and spruce sp., injury to Pinus murrayana. ponderosa. Dendrolene against two-lined chestnut-borer.	\$\begin{cases} 93, rev. 2 \\ 37, rev. 2 \\ 37, rev. 2 \\ 59 \\ 11, rev. \\ 80 \\ 51, rev. \\ 96 \\ \\ 87 \\ 72 \\ 75, rev. 2 \\ 24, rev. \\ 61 \\ 24, rev. \\ \\ 61 \\ 24, rev. \\ \\ \\ \\ \end{cases}	5 4 4 4 3 3 4 5 5 2 2 1 1 5 8 8 4 4 2 2 3 3 2 2 1 9	
Dactylis glomerata, food plant of Toxoptera graminum. Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Dandelion, food plant of Aphis gossppii. Däne," colloquial name for Blattella germanica. Daremma catalipa, same as Ceratomia catalipa. Dasyneura leguminicola. (See Clover-flower midge.) Datura, food plant of Aphis gossppii. Leptinotarsa decemlineata. probable food plant of Diabrotica 12-punctata. Davallia mooreana, food plant of Chionaspis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to pine and spruce. sp., injury to Pinus murrayana. ponderosa. Dendrolene against two-lined chestnut-borer. Dermanyssus gallina, general account. remedies.	\$\begin{cases} 93, rev. 2 \\ 37, rev. 2 \\ 37, rev. 2 \\ 59, \\ 59, \\ 80 \\ 80, \\ 87, \\ 24, rev. 2 \\ 24, rev. 2 \\ 29, \\ 92 \\ 92 \\ 92 \\ 92 \\ \ 92 \\ \ \ \	5 4 4 4 3 3 4 4 5 5 2 2 1 5 8 8 4 4 2 2 3 3 2 1 1 9 9 7 7 1 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata Daisy, ox-eye, food plant of Macrodactylus subspinosus. Dandelion, food plant of Aphis gossypti Dane," colloquial name for Blattella germanica Daremma catalpæ, same as Ceratomia catalpæ. Daspneura leguminicola. (See Clover-flower midge.) Datura, food plant of Aphis gossypti.  Leptinotarsa decemlineata. probable food plant of Diabrotica 12-punctata. Davallia mooreana, food plant of Chionaspis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to spine and spruce. sp., injury to Pinus murrayana ponderosa.  Dendrolene against two-lined chestmut-borer. Dermanyssus gallinæ, general account. remedies. Deutzias, food plants of Macrodactylus subspinosus.	\$\begin{cases} 93, rev. 2 \\ 37, rev. 2 \\ 37, rev. 2 \\ 37, rev. 2 \\ 11, rev. \\ 80 \\ 51, rev. \\ 96 \\ 87, rev. 2 \\ 24, rev. \\ 24, rev. \\ 92 \\ 11, rev. \\ 93, rev. 2 \\ 11, rev. \\ 92 \\ 11, rev. \\ 93, rev. 2 \\ 11, rev. \\ 93, rev. 2 \\ 11, rev. \\ 11, r	55 44 33 45 52 1 1 5 8 8 44 2 2 3 3 2 2 1 9 7 1 1 7	
Dactylis glomerata, food plant of Toxoptera graminum. Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Dandelion, food plant of Aphis gossypti. Däne," colloquial name for Blattella germanica. Daremma catalpæ, same as Ceratomia catalpæ. Dasyneura leguminicola. (See Clover-flower midge.) Datura, food plant of Aphis gossypti. Leptinotarsa decemlineata. probable food plant of Diabrotica 12-punctata Davallia mooreana, food plant of Chionaspis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to pine and spruce. sp., injury to Pinus murrayana. pondrosa. Dendrolene against two-lined chestnut-borer. Dermanyssus gallinæ, general account. remedies. Deutzias, food plants of Macrodactylus subspinosus.	\$\begin{cases} 93, rev. 2 \ 37, rev. 2 \ 37, rev. 2 \ 27, rev. 2 \ \ 80 \ 51, rev. \ 80 \ \ 87 \ \ 96 \ \ 87, rev. 2 \ 24, rev. \ 61 \ 24, rev. \ 92 \ 11, rev. \ 21,	54 44 33 45 52 21 58 44 22 33 22 11 77 1-33 42	
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Daisy, ox-eye, food plant of Macrodactylus subspinosus. Dandelion, food plant of A phis gossypii. 'Däne,'' colloquial name for Blattella germanica. Daremma catalpæ, same as Ceratomia catalpæ. Dasyneura leguminicola. (See Clover-flower midge.) Datura, food plant of A phis gossypii. Leptinotarsa decemiineata. probable food plant of Diabrotica 12-punctata. Devallia mooreana, food plant of Chionaspis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to pine and spruce. sp., injury to Pinus murrayana. ponderosa. Dendrolene against two-lined chestnut-borer. Dermanyssus gallinæ, general account. remedies. Deutzias, food plants of Macrodactylus subspinosus. Dewberry, food plant of Anthonomus signatus. Diabrotica 12-punctata general account.	\$\begin{cases} 93, rev. 2 \\ 37, rev. 2 \\ 37, rev. 2 \\ 59 \end{cases} 11, rev. \\ 80 \\ 51, rev. \\ 96 \\ 80 \\ 87, rev. 2 \\ 24, rev. \\ 92 \\ 11, rev. \\ 21, rev. \\ 21, rev. \\ 59 \end{cases} \end{cases} 11, rev. \\ 21, rev. \\ 21, rev. \\ 59 \end{cases} \end{cases} \end{cases} \text{11}, rev. \\ 21, rev. \\ 59 \end{cases} \text{12}, rev. \\ 21, rev. \\ 59 \end{cases} \text{13}, rev. \\ 21, rev. \\ 59 \end{cases} \text{14}, rev. \\ 21, rev. \\ 59 \end{cases} \text{15}, rev. \\ 21, rev. \\ 21, rev. \\ 59 \end{cases} \text{15}, rev. \\ 21, rev. \\ 21, rev. \\ 59 \end{cases} \text{15}, rev. \\ 21,	5 4 4 3 3 4 5 2 1 1 5 8 4 2 2 3 2 1 1 1 3 4 5 7 1 1 1 3 4 1 1 3 1 1 3 1 3 4 1 1 3 1 3 1	1-
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Danisy, ox-eye, food plant of Macrodactylus subspinosus. Dandelion, food plant of Aphis gossypti. Däne," colloquial name for Blattella germanica. Daremma catalipæ, same as Ceratomia catalipæ. Dasyneura leguminicola. (See Clover-flower midge.) Datura, food plant of Aphis gossypti. Leptinotarsa decemlineata. probable food plant of Diabrotica 12-punctata. Davallia mooreana, food plant of Chionaspis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to pine and sprue. sp., injury to Pinus murrayana. ponderosa. Dendrolene against two-lined chestnut-borer. Dermanyssus gallinæ, general account. remedies. Deutzias, food plants of Macrodactylus subspinosus. Dewberry, food plants of Macrodactylus subspinosus.	\$\begin{cases} 93, rev. 2 \\ 37, rev. 2 \\ 37, rev. 2 \\ 27, rev. 80 \\ 51, rev. \\ 80 \\ 51, rev. 2 \\ 75, rev. 2 \\ 24, rev. \\ 61 \\ 24, rev. \\ 21, rev. \\ 25, 559 \\ 599 \\ 25, \end{cases}	54 44 33 45 52 21 58 44 22 33 22 11 77 1-33 42	1-
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Dalsy, ox-eye, food plant of Macrodactylus subspinosus. Dandelion, food plant of Aphis gossypii 'Diane,'' colloquial name for Blattella germanica. Daremma catalpæ, same as Ceratomia catalpæ. Dasyneura leguminicola. (See Clover-flower midge.) Datura, food plant of Aphis gossypii.  Leptinotarsa decemiineata. probable food plant of Diabrotica 12-punctata.  Davallia mooreana, food plant of Chionospis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to pine and spruce. sp., injury to Pinus murrayana. ponderosa.  Dendrolene against two-lined chestnut-borer. Dermanyssus gallinæ, general account. remedies. Deutzias, food plant of Anthonomus signatus. Diabrotica 12-punctata, general account. remedies. longicornis, general account. remedies.	\$\begin{array}{c} 93, rev. 2 \\ 37, rev. 2 \\ 37, rev. 2 \\ 37, rev. 2 \\ 61 \\ 61 \\ 61 \\ 24, rev. 2 \\ 11, rev. 2 \\ 24, rev. 2 \\ 24, rev. 2 \\ 25, rev. 3 \\ 25, rev.	5 4 4 3 4 5 5 2 1 1 5 8 8 4 4 2 2 1 1 9 9 7 1 1 3 3 4 5 1 1 1 1 3 1 1 3 1 1 1 1 3 1 1 1 1 1 1	1-
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Dahlia, food plant of Diabrotica 12-punctata. Daisy, ox-eye, food plant of Macrodactylus subspinosus. Dandelion, food plant of Aphis gossypii. 'Däne,'' colloquial name for Blattella germanica. Daremma catalpæ, same as Ceratomia catalpæ. Dasyneura leguminicola. (See Clover-flower midge.) Datura, food plant of Aphis gossypii. Leptinotarsa decemlineata. probable food plant of Diabrotica 12-punctata. Davallia mooreana, food plant of Chionaspis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to pine and spruce. sp., injury to Pinus murrayana ponderosa. Dendrolene against two-lined chestnut-borer. Dermanyssus gallinæ, general account. remedies. Deutzias, food plants of Macrodactylus subspinosus. Diabrotica 12-punctata, general account. remedies. longicornis, general account. remedies. soror, pest in Pacific States.	\$\begin{cases} 93, rev. 2 \ 37, rev. 2 \ 37, rev. 2 \ 59 \ 59 \ 59 \ 59 \ 59 \ 59 \ 59 \	5 4 4 4 3 3 4 4 5 5 2 1 1 5 8 8 4 4 2 2 3 3 2 2 1 1 9 9 7 7 1 3 4 4 2 2 1 , 2 - 5 5 1 , 6 - 8 8 3 3	1-
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Daisy, ox-eye, food plant of Macrodactylus subspinosus. Dandelion, food plant of Aphis gossypii. 'Cbäne,'' colloquial name for Blattella germanica. Daremma catalpæ, same as Ceratomia catalpæ. Dasyneura leguminicola. (See Clover-flower midge.) Datura, food plant of Aphis gossypii.  Leptinotarsa decemiineata. probable food plant of Diabrotica 12-punctata.  Davallia mooreana, food plant of Chionaspis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to pine and spruce. sp., injury to Pinus murrayana. ponderosa. Dendrolene against two-lined chestnut-borer. Dermanyssus gallinæ, general account. remedies. Deutzias, food plants of Macrodactylus subspinosus. Deutzias, food plants of Macrodactylus subspinosus. Diabrotica 12-punctata, general account. remedies. longicornis, general account. remedies. soror, pest in Pacific States. vittata, carrier of fungous diseases of plants.	\$\begin{cases} \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qqquad \qqqq\qqq\qqq\qqq\qqq\qqq\qqq\qqq\qqq\	5 4 4 3 4 5 2 1 1 5 8 4 4 2 2 3 3 2 1 1 9 7 1 1 3 4 4 5 1 1 1 1 3 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata Daisy, ox-eye, food plant of Macrodactylus subspinosus. Dandelion, food plant of A phis gossypii. "Dâne," colloquial name for Blattella germanica. Daremma catalpæ, same as Ceratomia catalpæ. Dasyneura leguminicola. (See Clover-flower midge.) Datura, food plant of A phis gossypii. Leptinotarsa decemiineata. probable food plant of Diabrotica 12-punctata.  Darallia mooreana, food plant of Chionaspis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to pine and spruce. sp., injury to Pinus murrayana ponderosa. Dendrolene against two-lined chestnut-borer. Dermanyssus gallinæ, general account. remedies. Deutzias, food plant of Anthonomus signatus. Diabrotica 12-punctata, general account. remedies. soror, pest in Pacific States. vittata, carrier of fungous diseases of plants circular enneny of cucurbits.	\$\begin{array}{c} \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qqquad \qqqq \qqqqq \qqqq \qqqq \qqqq \qqqq \qqqq \qqqqq \qqqqq \qqqqq \qqqqq \qqqqq \qqqqqq	5 4 4 4 3 3 4 4 5 5 2 1 1 5 8 8 4 4 2 2 3 3 2 2 1 1 9 9 7 7 1 3 4 4 2 2 1 , 2 - 5 5 1 , 6 - 8 8 3 3	
Dactylis glomerata, food plant of Toxoptera graminum.  Dactylopius adonidum on grape. remedy. Dahlia, food plant of Diabrotica 12-punctata. Daisy, ox-eye, food plant of Macrodactylus subspinosus. Dandelion, food plant of A phis gossypii. "Dainerma catalpæ, same as Ceratomia catalpæ. Dasyneura leguminicola. (See Clover-flower midge.) Datura, food plant of A phis gossypii.  Leptinotarsa decemiineata. probable food plant of I Diabrotica 12-punctata.  Davallia mooreana, food plant of Chionaspis sp. Deinocerites cancer, distinctive characters of larva. Delaware, requirements relative to shipment of nursery stock. Dendroctonus frontalis, destructive to pine and spruce. sp., injury to Pinus murrayana. ponderosa. Dendrolene against two-lined chestnut-borer. Dermanyssus gallinæ, general account. remedies. Deutzias, food plants of Macrodactylus subspinosus. Deuberry, food plants of Anthonomus signatus. Diabrotica 12-punctata, general account. remedies. longicornis, general account. remedies. soror, pest in Pacific States. vittata, carrier of fungous diseases of plants.	\$\begin{array}{c} \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qqquad \qqqq \qqqqq \qqqq \qqqq \qqqq \qqqq \qqqq \qqqqq \qqqqq \qqqqq \qqqqq \qqqqq \qqqqqq	5 4 4 3 3 4 4 5 5 2 1 1 5 8 8 4 4 2 2 3 3 2 2 1 1 9 9 7 7 1 3 3 4 4 2 5 5 1,6 8 8 8 3 7 1 1 8 8 8 8 7 1 1 8 8 8 8 7 1 1 8 8 8 8	1

	Circular No.	Page.	Figure.
Diatræa saccharalis [larger corn stalk-borer], general account	16	1-3	1-3
remedies. [sugar-cane borer], distribution. Dibrachys boucheanus, secondary parasite of Thyridopteryx ephemeræ-	16 16	3	• • • • • • • • • • • • • • • • • • • •
formis.  Dieffenhachia jenmani furnication with bydrovania acid gas	97 37, rev. 2	7-8	10
Diglochis omnivorus, parasite of Malacosoma americana	98	3 5	
Dioracnys obucheanus, secondary parasite of Thyriaopteryx ephemeræ- formis.  Dieffenbachia jenmani, fumigation with hydrocyanic-acid gas.  Dieffenbachia jenmani, fumigation with hydrocyanic-acid gas.  Diplochis omnivorus, parasite of Malacosoma americana.  Diodia, food plant of Aphis gossypii.  Diplodus luridus, enemy of Malacosoma americana.  Diplosis. (See Contarinia.)  tritici. (See Contarinia tritici.)  Disease, bacterial wilt, of curcubits, cotton, and cowpeas. (See  Bacillus tracheiphilus and Neocosmospora vasinfecta.)  fungous, of grasshoppers.	80 98	5 6	
Disease, bacterial wilt, of curcubits, cotton, and cowpeas. (See Bacillus trachetphilus and Neocosmospora vasinfecta.)			
fungous, of grasshoppers. leaf-spot, of catalpa. (See Phyllosticta catalpæ.) of Malacosoma americana.  Diseases, human, carriage by Musca domestica of brood of bees, circular.	84	5–7	6-7
of Malacosoma americana	98 71, rev.	6 3	
of brood of bees, circular	71, 167.		
remedies. chickens attributed to presence of mites and lice. induced by lice. plants, carriage by <i>Diabrotica longicornis</i> and striped cucumber beetle.	79 92	3-5	
induced by lice.	92	5	
striped cucumber beetle	59	7	
transmission by Anasa tristis	39, rev. 2 31, rev. 2	2 2	
Discontaing agraling boot of Marmis albigage	69, rev.	3	
Districtly spicata, food plant of Toxoptera graminum.  Ditropinotus aureoviridis, parasite of Isosoma tritici.	93, rev. 66, rev.	5 4	
Dock. (See Rumex.)			
Documents, destruction by Termes flavipes	50, rev. 13	3-4	
Dogwood. (See Cornus.) DORSETT, P. H., and Woods, Albert F., circular, "The Use of Hydrocyanic Acid Gas for Fumigating Greenhouses and Cold	99		
Frames"	37, rev.	1-10	1-3
Doryphora 10-lineata=Leptinotarsa decemlineata "Drill worm," colloquial name for Diabrotica 12-punctata	87 59	$\frac{1}{2}$	
Drosophila ampelophila in houses	71, rev.	$3, \bar{5}$	7
	79	3	
Drying against chestnut weevils  Duck, enemy of Leptinotarsa decembineata	99 87	10–11 12	
Dust against pear slug	31, rev. 2	6	
Drying against chestnut weevils. Drying against chestnut weevils. Duck, enemy of Leptinotarsa decemlineata. Dust against pear slug. striped cucumber beetle. and sulphur against cotton red spider. DYAR, HARRISON G., circular, "Key to the Known Larvæ of the Mosquitoes of the United States".	65	4	
Mosquitoes of the United States"	72	1-6	1
	CO		
Earthworms mistaken for Mermis albicans.  Echinocystis lobata, food plant of Diabrotica vittata.  Melittia satyriniformis.	62, rev. 2 31, rev. 2 38, rev. 2	5 2 3	
Ectobia germanica. (See Blattella germanica.) Eggplant, food plant of Aleyrodes vaporariarum	57	2	
Leptinotarsa decemlineata  Eichhornia speciosa, fumigation with hydrocyanic-acid gas	87	8 3	
Elder, food plant of Thyridopteryx ephemeræformis	37, rev. 2 97	4	
Elder, food plant of Thyridopteryx ephemeræformis.  Elm, food plant of Bryobia pratensis.  Galerucella luteola.	19, ed. 2 8, rev.	2 1-4	
Malacosoma americana	98	2	
Pulvinaria innumerabilis	29, rev.	2 2	
Scolytus rugulosus. Thyridopteryx ephemeræformis. leaf-beetle, circular.	97	4	
	8, rev. 15	1-6 3,4	1
remediespossible food plant of Chrysobothris femorata	8, rev. 3 32, rev. 3	4-6 10	
Elymps, food plant of Mayetiola destructor or related species	70	7	
striatus, food plant of Toxoptera graminum.  Empusa aphidis, fungous disease of Macrosiphum pisi.	93, rev. 43, ed. 2	5	
muscx, fungous disease of Musca domestica	35	8	
Encyrtus sp., parasite of Eriocampoides limacina.	71, rev. 26	9	
Entedon epigonus, parasite of (Cecidomyia) Mayetiola destructor	12, rev.	3 11	
Evania appendigaster	51, rev.	11	
pular	76	1-32	
Epicauta vittata, an "old-fashioned potato bug". Epilampra notabilis, predaceous upon herbivorous larvæ. Eragrostis, possible food plant of Toxoptera graminum	51, rev.	3 2	3
Eragrostis, possible food plant of Toxoptera graminum  Eriocampoides limacina, circular	85 26	1-7	1-4
remedies	26	6-7	

	Circular No.	Page.	Figure.
·			
Eulecanium nigrofasciatum, circular	88	1-4	1–3
remedies.  persicx, confounded with Eulecanium nigrofasciatum  prunastri.	88 88	3-4	
nrunastri.	88	3	
Eunotus lividus, parasite of Pulvinaria innumerabilis	64	4	
Eupelmus allynii, parasite of Isosoma tritici	66, rev.	4	
Mayetiola destructor	39, rev. 2	12 4	12-13
spp. parasites of Scolutus rugulosus	29, rev. 2	6	
Eupeodes volucris, enemy of Aphis gossypii	80	6	
reduvii, parasite of Anasa trisits.  spp., parasites of Scolylus rugulosus. Eupeodes volucris, enemy of Aphis gossypii. Euphorocera claripernis, parasite of Ceratomia catalpæ. Euproctis chrysorrhæa, character, conditions in New England, July,	96	5	2
control by collecting and burning winter	58	10–12	
webs	58 29, rev.		
crassineum, parasite of Scolutus rugulosus	29, rev.		
crassineum, parasite of Scolytus rugulosus. Euschistus variolarius, enemy of Leptinotarsa decemlineata. Eutelus flavipes, parasite of Cheilosia alaskensis.	87		
Eutelus flavipes, parasite of Cheilosia alaskensis	61		
	68	1-5	
remedles.	68 68	2-5	
Evania appendiaaster, host of Entedon hagenowi	51, rev.	11	
remedies.  tritici, the wheat thrips.  Evania appendigaster, host of Entedon hagenowi parasite of cockroaches	51, rev.	111	
excrement (see also manure).			
human, breeding place of Musca domestica	71, rev. 80	. 6	
Exorista vulgaris, parasite of Pontia rapx	60	6	
			1
<b>F.</b>			
	[ 14	7-8	
	18	7-8	
Tall destruction of cotton stalks against boll weevil	27 33	6-7	
Ÿ	56	4-5	
	95, rev.		
treatment against American and European foul brood	79	4	
Feathers, food of Attagenus piceus	36, rev.	4	
$Tineola\ biselliella$	36, rev. 36, rev.		
Felting, food of Trichophaga tanetzella	36, rev.	5, 5	
Pelting, food of <i>Trichophaga tapetzella</i> Perns, fumigation with hydrocyanic-acid gas	37, rev. 2	2	
ertilizers (see also Manure).	10		
àgainst Hessian flyjoint-worm	66, rev.	3 5	
commercial, against corn root-aphis	86	12	
mineral, against seed-corn maggot	63, ed. 2	3	
organic, inducing root-maggot infestation	63, ed. 2	3	
Ficus elastica, fumigation with hydrocyanic-acid gas	37, rev. 2	3	
bark maggot, lowland. (See Cheilosia hoodianus.) grand. (See Fir, lowland.)	61	9,10	
lowland, nost tree of Cheilosia hoodianus	61	8-9,10	
Hylesinus n. sp., and Hylesinus granu- latus	61	9	
white. (See Fir, lowland.)			
Fire-brat," colloquial name for Lepisma domestica	49	3	
Fire-brat," colloquial name for Lepisma domestica. Fireplaces, infestation by Lepisma domestica Fish bait, larvæ of Ceratomia catalpæ used therefor	49	3	
Fish bait, larvæ of Ceratomia catalpæ used therefor	96	4	
dried, food of cockroaches	51, rev. 24, rev.	2 7	
soap. (See Soap, fish-oil, and Soap, whale-oil.) Flea, cat and dog. (See Pulex serraticeps.) human. (See Pulex irritans.)	24,161.	,	
human. (See Pulex irritans.)			
fleas, general account	46, rev.	3-4 1,2	
hydrocyanic-acid gas as remedy	46, rev. 13	1,2	
Flies feeding on honeydew from aphides	80	i	
	ſ 35	8	
	71, rev.	3	
house, as carriers of disease.	35 71, rev.	1-8 1-10	1-6
circulars.		2	1-:
circulars	46. rev.		
circularshydrocyanic-acid gas as remedy	46, rev. 35	5-8	
circulars. hydrocyanic-acid gas as remedy. remedies	46, rev. 35 71, rev.	5-8 5-9	
circulars. hydrocyanic-acid gas as remedy. remedies. species occurring in houses.	46, rev. 35 71, rev. 35 71, rev.	5–8 5–9 1–3 1–6	
circulars. hydrocyanic-acid gas as remedy. remedies	46, rev. 35 71, rev. 35 71, rev. 84	5–8 5–9 1–3 1–6	

	Circular No.	Page.	Figure.
Flour and sulphur against cotton red spider. Fly, bluebottie. (See Calliphora crythrocephala and Lucilia czsar.)	65	4	
cluster. (See Pollenia rudis.) fruit. (See Drosophila ampelophila.)			
green-bottle. (See Lucilia casar.)			
Hessian. (See Cecidomyia destructor and Mayetiola destructor.) house. (See Musca domestica.)			
meat. (See Calliphora erythrocephala.)		_	
paper, sticky, against house fly	35 71, rev.	5	
small bluebottle. (See Phormia terrænovæ.) house. (See Homalomyia canicularis.)			
stable. (See Cyrtoneura stabulans, Muscina stabulans, and		,	
Stomoxys calcitrans.) typhoid. (See Musca domestica.)			
typhoid. (See Musca domestica.) window. (See Scenopinus fenestralis.)			
Food, infestation by cockroaches	51, rev. 34, rev.	4-5	
Levisma saccharina	49	1	
Forage plants, food of <i>Bryobia pratensis</i> Forestalling outbreaks of grasshoppers Forest trees, food plants of <i>Macrodactylus subspinosus</i> .	19, ed. 2 84	10	
Forest trees, food plants of Macrodactylus subspinosusinjury by periodical cicada	11, rev. 30	$\frac{2}{1}$	
Formalin against powder-post heetles	55	4	
Formica fusca, enemy of Anthonomus signatus.	21, rev. 94		-,
Formica fusca, enemy of Anthonomus signatus. Foul brood, American, cause, circular description not the foul brood of Cheshire and Cheyne	79	1-2	
treatment	79		
European, description	79	2	
treatment of Cheshire and Cheyne, not American foul brood	79 94	3-4	
Fowle (can also Poultry and Chicken)	92		
enemies of boll weevil.	27	5-6	
diseases induced by lice. enemies of boll weevil. grasshoppers. Leptinotarsa decemlineata.	84 87	4 12	
yellow. (See Ixophorus glaucus.)	. 01	12	
France, laws relative to importation of American plants, trees, and fruits.	41	2	
"Franzosen," colloquial name for Blattella germanica.  Frog, tree, enemy of cockroaches.	51, rev. 51, rev.	2 11	
Frontina frenchii, parasite of Ceratomia catalpa	96	5	
Frost. (See Temperature and Weather.)	98	6	
Fruit flies, quarantine regulations of New England against intro-	41	4	
duction fly. (See Drosophila ampelophila.)	41	-	
trees, food plants of Bryobia pratensis. injury by buffalo tree-hopper.	19, ed. 2 23	1, 2, 3, 4 1-3	
protection against periodical cicada	22	1	
	22 30		
young, injury by periodical cicada	44	1-2	
	74		
Fruits, food of cockroaches garden, food plants of Macrodactylus subspinosus.	51, rev. 11, rev.	3 2	
	28	1,2	
Filensia, 1000 Diant of Alegrodes raporariorum	57 37, rev. 2	2 3	
fumigation with hydrocyanic-acid gas Fumago salicina, fungus following work of Fsylla pyricola. Fumigation. (See Carbon bisulphid, Hydrocyanic-acid gas, and	7	2	
under Tobacco.) Fungi, food of <i>Diabrotica longicornis</i> Fungous disease, believed to have caused disappearance of <i>Dendroc</i> -	59	7	
tonus frontalis	24, rev.	1	C P
of grasshoppers  Macrosiphum pisi	43, ed. 2	<b>5–7</b>	0-7
Musca domestica. (See Empusa muscx.) Phytonomus punctatus	69	9	
plants, carriage by Diabrotica longicornis	. 59	7	
striped cucumber beetle  Pollenia rudis	59 71, rev.	. 7	
Fungus, smoky. (See Fumago salicina.) Fur, food of Tinea pellionella. Tineola biselliella.	36, rev.	3 4	
Trichophaga tapetzella	36, rev.	5	
Furniture, destruction by Termes flavipes.  plush-covered, reported injury by Lepisma saccharina	50, rev.	4,5	
protection from injury by white ants		8	

	Circular No.	Page.	Figure.
G.			
Galeruca xanthomelæna. (See Elm leaf-beetle.) Galerucella luteola. (See Elm leaf-beetle.)			•
Garbage, breeding place of Musca domestica	35	5	
	71, rev. 63, ed. 2	3 5	
Gas lime against cabbage maggot. root form of woolly aphis.	20, rev.	4	
impracticable against house flies	35 71, rev.	6	
tar against peach-tree borer	54	4	
	3	8-9	
Gasoline against powder-post beetles.  Geophilus spp., mistaken for Mermis albicans.  Georgia, requirements relative to shipment of nursery stock.  Geranium, food plant of Aleyrodes vaporariorum.  Termes flavipes.  Germany, laws relative to importation of American plants, trees, and fruits.	62, rev.	4 5	
Georgia, requirements relative to shipment of nursery stock	75, rev. 2	2-3	
Geranium, food plant of Aleyrodes vaporariorum	.57	2	
Cormany laws relative to importation of American plants trees	50	6	
and fruits	41	2-3	
Giant whips. (See Spirochæta apis.) Gipsy moth. (See Porthetria dispar.) GRAULT, A. A., JOHNSON, FRED, and, circular, "The Plum Curculio (Conotrachelus nenuphar Herbst)". Girdling against fruit-tree bark-beetle.			
Greatury A. A. Johnson Epen and circular "The Plum Curculia			
(Conotrachelus nenuphar Herbst)".	73	1-10	1-5
Girdling against fruit-tree bark-beetle.	29, rev.	7	
at certain times to prevent pinhole injury in cypress  Goldenrod flowers, frequented by adults of Cyllene robiniæ	82 83	3	
food plant of Diabrotica 12-punctata		3	
longicornis	59	7	
Vittata	02	7	6
Goniodes dissimilis, parasite of chickens "Gooseberry bugs," colloquial name for chiggers food plant of Pulvinaria innumerabilis.	92	7	5
"Gooseberry bugs," colloquial name for chiggers	77	4	
Gourd as trap plant against striped cucumber beetle	31, rev. 2	2 6	
food plant of Melittia satyriniformis	38, rev. 2	2	
Grabhamia discolor, distinctive characters of larva jamaicensis, distinctive characters of larva.	72	3	
Grain anhis European (See Sinhocorung avense)	72	3	
Grain aphis, European. (See Siphocoryne avenæ.) spring. (See Toxoptera graminum.)			
moth Angoumois (Soo Sitotroga agrealella)	40		
weevils, hydrocyanic-acid gas as remedy.  Grains, food plants of Camnula pellucida.	46, rev. 84	2	
small, food plants of Leucania unipuncta	4	1	
Granaries, fumigation with hydrocyanic-acid gas	46, rev.	2 2	
Granary beetles, hydrocyanic-acid gas as remedy.  Grape, food plant of Dactylopius adonidum.	46, rev. 37, rev. 2	4	
Diabrotica 12-punctata	59	3	
Macrodactyluŝ subspinosus Pulvinaria innumerabilis	11, rev. 64	1-2	
Phylloxera, resin washes as insecticide.	1	5	
Granes under glass, furnigation with hydrocyanic-acid gas	. 37 rev	4	
Grass, beard. (See Polypogon monspeliensis.) blue, food plant of Leucania unipuncta stem, food plant of Bryobia pratensis.	4	1	
stem, food plant of Bryobia pratensis	19, ed. 2	2	
couch, food plant of Toxoptera graminum		4	
foxtail, food plant of Aphis maidi-radicis.	86	6	
gama. (See Tripsacum dactyloides.) orchard. (See Dactylis glomerata.) sesame. (See Tripsacum dactyloides.)			
sesame. (See Tripsacum dactyloides.)			
squirrel-tail. (See Hordeum jubatum.) Grasses, food of Termes flavines	50, rev.	6	
Grasses, food of Termes flavines  plants of Bryobia pratensis.  Camnula pellucida.  Leucania unipuncta.	19	2, 3, 4	
Camnula pellucida	84	1	
Macrodactylus subspinosus,	11, rev.	2	
Mayetiola destructor or related species	70	7.	
Grasshopper, differential. (See Melanoplus differentialis.) pellucid. (See Camnula pellucida.)			
problem and alfalfa culture, circular	84	1-10	1-8
Rocky Mountain. (See Melanoplus spretus.) two-striped. (See Melanoplus bivittatus.)			
two-striped. (See Melanoplus bivittatus.) yellow-winged. (See Camnula pellucida.)			
Grasshoppers, remedies.	84	6-10	8
Grazing against boll weevil	56 05 TOW	4 6	
	95, rev.	9	
"Green arsenoid" against plum curculio.  Green-bottle fly. (See Lucilia cæsar.)			
"Green bug," colloquial name for Toxoptera graminum	85 O3 TOV	1	
	\ .93, rev.	1	
Greenhouse white fly. (See Aleyrodes vaporariorum.) Greenhouses and cold frames, fumigation with hydrocyanic-acid gas,	07 0	4 44	1.0
circular infestation by Aleyrodes vaporariorum	37, rev. 2 57	1-11	1-3
into two of 21 oground to portation and	01		

	Circular No.	Page.	Figure.
Greenhouses, infestation by Diabrotica vittata	31, rev. 2	2	
Leptocoris trivittatus	28	3	
Grosbeak, rose-breasted, enemy of Leptinotarsa decemlineata	50, rev.	6	
Ground beetle, fiery. (See Calosoma calidum.)	87	11	
slender seed-corn. (See Clivina impressifrons.)			
beetles, enemies of Leucania unipuncta	4	4, 5	
Groundsel. (See Senecio.)		10	
Grouse, ruffed, enemy of Leptinotarsa decemlineata "Grubby cattle," meaning of term.	87 25	12	
"Grub," colloquial name for Hypoderma lineata.	25	1	
Grubs, white, kerosene emulsion as insecticide.	1	5	
Gum, injury by Platypus compositus	82	3	
Gunpowder against cockroaches Gypsum and kerosene against squash bug	51, rev.	5	
turpentine against squash bug	39, rev. 2 39, rev. 2	5	
Gypsy moth. (See Porthetria dispar.)	00,101.2	Ĭ	
н.			
Habrocytus thyridopterigis, hyperparasite of Thyridopteryr ephemer-			
æformis	97	8	11
probable parasite of Allocota thyridopterigis	. 019	8	
Hackberry, food plant of Pulvinaria innumerabilis	97 64	2	
Hadronotus anasæ, parasite of Anasa tristis	39, rev. 2	4	
Hair, food of Tineola biselliella	36	4	
worm, cabbage (see also Mermis albicans).	60	3	
positively not poisonous	62, rev.		
Hand destruction against house centipede	48	4	
method against plum curculio.	73	10	
methods against seed-corn maggot. squash-vine borer	63, ed. 2	\ 5-6	
picking against apple-tree tent caterpillar	38, rev. 2	7	
	97	8-9	
cabbage maggot	63, ed. 2	4	
cabbage maggot. catalpa sphinx imported cabbage worm.	96	6 8	
periodical cicada	60   89	4	
rose-chafer	11, rev.	3-4	
Squash bug.  Harpalus caliginosus, enemy of Leptinotarsa decemlineata.	39, rev. 2	4	
Harvest mites or "chiggers," circular	87	10 1-6	1-3
remedies	77	5-6	1-0
Hawaii, requirements relative to shipment of nursery stock	75, rev. 2	3	
Haw, food plant of Conotrachelus nenuphar	73	4 4	
Hawk, sparrow, enemy of grasshoppers. Swainson, enemy of grasshoppers.	84 84	4	
Hawthorn, food plant of Pulvinaria innumerabilis.	64	2	
Scolytus rugulosus	29, rev.	2	
Hazel, food plant of Balaninus obtusus	99	15	
Hazelnut weevil. (See Balaninus obtusus.) Heat. (See Temperature and Weather.)			
Heating against chestnut weevils.  powder-post beetles.  "Heel fly," colloquial name for Hypoderma lineata.	99	11	
powder-post beetles.	55		
"Heel fly," colloquial name for Hypoderma lineata	25		
Heliothis obsoleta, Paris green as remedy.  Heliotrope, food plant of Aleyrodes vaporariorum.	68 57	3-4	
Heliebore against imported cabbage worm  pear slug  ineffective against rose-chafer.  Hemerobius gossypii, enemy of Aphis gossypii.  Hemerosyma levestiyan (Sos Tussock noth white-marked)	60	8	
pear slug	26	7	
ineffective against rose-chafer	11, rev.	3	
Hemerocampa leucostigma. (See Tussock moth, white-marked.)	80	0	
Hemiteles mesochoridis, parasite of A panteles congregatus	96	5	
scolyti, parasite of Scolytus rugulosus	29, rev.	6	
thyridopterigis=Allocota thyridopterigis	97	6	
Hemlock bark beetle. (See Hylesinus n. sp.) maggot. (See Cheilosia alaskensis.)			
Western, "Black check" injury, circular	61	1-10	1-5
host tree of Cheilosia alaskensis	61	2-8	1,3-5
Hylesinus n. sp	61	- 2,3	3
Henbane. (See Hyoscyamus.) Hen louse. (See Menopon pallidum.)	į		
manure, ineffective against strawberry weevil	21, rev.	7	
Herbariums, reported infestation by Tineola biselliella	36, rev.	4	
Hessian fly. (See Cecidomyia destructor and Mayetiola destructor.)			
Heteropus ventricosus. (See Pediculoides ventricosus.) Heydenia unica, perasite of Scolytus rugulosus	29, rev.	6	
Hibiscus. (See Mallo and Mallows.)	Do, ICV.	0	
Hickory, food plant of Chrysobothris femorata	32, rev. 3	9	
Hippodamia convergens, enemy of Aphis gossypii	80	6,7	3, a, b, c
Leptinotarsa decemlineata	87 l	10	4

	Circular No.	Page.	Figure.
Hippodamia convergens, enemy of Macrosiphum pisi	43, ed. 2	5	
Toxoptera graminumglacialis, enemy of Leptinotarsa decemlineata	85 87	10	3
Macrosiphum pisi	43		
parenthesis, enemy of Toxoptera graminum	85	5	
13-punctatá, enemy of A phis gosšypii	80 87	6 10	
Macrosiphum pisi	43, ed. 2		
Ioffmannia refulgens, fumigation with hydrocyanic-acid gas	37, rev. 2	3	
Jonalodisca coggulata glassy-winged sharpshooter mistaken for	6	2,3	3
boll weevil	27	5	
Iomalomyia brevis in houses	71. rev.	2.4	(
canicularis in houses	$ \begin{cases} 35 \\ 71, \text{rev.} \end{cases} $	2	
Tomoporus chalcidiphagus, parasite of Isosoma tritici	66, rev.	4	
Innovedow from corn root applic	86	8	
Prevent 'White Veins' in Wrapper Tobacco (Euthrips nicotiana)	69	1 7	
Hinds)"	68 80	1-5 5	
Hinds)". Iop, food plant of Aphis gossypii. Phorodon humuli.	2	3-4	
Topkins, A. D., circular, "Pinhole Injury to Girdled Cypress in			
the South Atlantic and Gulf Štates" "Powder-Post Injury to Seasoned Wood	82	1-4	1
Products''	55	1-5	1
Products". "The Locust Borer and Methods for its Control".			
Control"	83	1-8	1-4
Toppordozor against grasshoppers	90 84	1-8 8-9	1-6
"The White-Pine Weevil".  Hopperdozer against grasshoppers. yellow-winged locust.	53	2-3	
		5	
jubatum, food plant of Toxoptera graminum. murinum, food plant of Toxoptera graminum. pusillum, food plant of Toxoptera graminum. Horismenus microgastri, parasite of Microplitis (Apanteles) catalpæ.	93, rev.	5	
murinum, 1000 plant of Torontera graminum.	93, rev.	4 5	
Horismenus microgastri, parasite of Microplitis (Apanteles) catalpx.	96	5	
Horse-enesting, 1000 plant of Chrysopothers femorale	32, rev. 3	9	
Horse droppings, salt, and Paris green against grasshoppers	84	9 2	
host of Dermanyssus gallinæ mint. (See Monarda fistulosa.)	92	2	
nettle, food plant of Diabrotica 12-punctata.  Horseradish, food plant of Pontia rapæ.	59 60	3 4	
House continede (See Scutigera forcens)			
cleaning against carpet beetle.	5, rev.	3-4	
clothes moths	36, rev.		
fly. (See Musca domestica.)	1	-	
small. (See Homalomyia canicularis.)	40 7 0		
Houses, infestation by Bryobia pratensis	19, ed. 2 47, rev.		
Cimex lectulariussp., from swallow	47, rev.		
cockroaches	51, rev.	1-14	
flies, circulars	35	1-8	1-6
Lepisma saccharina	71, rev.	1-10 1-4	
Leptocoris trivittatus	28	1.3	
Monomorium minutum	34, rev.	1.3	
pharaonis	34, rev.	1,2	1.0
Scutigera forceps, circular	48 34, rev.	1-4 $1, 3-4$	
Termes flavipes	50, rev.	1.0 1	
Howard, L. O., circular, "An Important Enemy to Fruit Trees.			
The San Jose Scale; its Appearance in the Eastern United States; Measures to			
be Taken to Prevent its Spread and to			
Destroy it''. "General Work against Insects which De-	. 3	1-10	1-8
"General Work against Insects which De-	1		,
foliate Shade Trees in Cities and Towns"	15 35	1-4 1-8	1-4
"House Flies (Musca domestica et al.)"	71, rev.	1-10	1-9
"How to Distinguish the Different Mos-	( 12)		
quitoes of North America"	40	1-7	1-3
"Hydrocyanic-acid Gas against House- hold Insects"	46, rev.	1-7	
hold Insects". "Mosquitoes and Fleas"	13	1-4	
"Regulations of Foreign Governments			
Regarding Importation of American Plants, Trees, and Fruits"		3 4	
	41	1-4	
"Remedial Work against the Mariaan		1.6	
"Remedial Work against the Mexican Cotton-boll Weevil"	33	1-6	
Cotton-boll Weevil'' "The Army Worm (Leucania unipuncta	33		
Cotton-boll Weevil''	33	1-5	1-3

	Circular No.	Page.	Figure.
HOWARD, L. O., circular "The Carpet Beetle or Buffalo Moth"			
(Anthrenus scrophulariæ L.)" "The Harlequin Cabbage Bug or Calico	5, rev.	1-4	1
"The Harlequin Cabbage Bug or Calico	10	1.0	F13
Back (Murgantia histrionica Hahn)" "The Larger Corn Stalk-horer (Diatræa	10	1–2	[1]
"The Larger Corn Stalk-borer (Diatræa saccharalis Fab.)"	16	1-3	1-3
"The Mexican Cotton-boll Weevil (An-	6	1-5	1-3
thonomus grandis Boh.)"	14 18	1-8 1-8	1-8 1-5
"The Mexican Cotton-boll Weevil in	10	1-0	1
1897''	27	1-7	
and Chittenden, F. H., circular, "The Bagworm (Thyridop-	1		
terux ephemer-	1		
æformis Haw.)" "The Catalpa		4 40	2.11
Haw.)	97	1–10	1-11
Sphinx ( Cera-			
tomia catalpæ Bdv.)"			
Bdv.)"	96	1-7	1-2
TUNTER, W. D., circular, "Note on the occurrence of the North American fever tick on sheep"	91	1-3	
"The Most Important Step in the Control	01	1-0	
"The Most Important Step in the Control of the Boll Weevil"	95, rev.	1-8	
"The Most Important Step in the Cui-			
tural System of Controlling the Boll Weevil"	56	1-7	
"The Periodical Cicada in 1902"	44	1-4	1-2
Hydrangea, food plant of Aphis gossypii	80	5	
Hydrocyanic-acid gas against bedbugs.	47, rev. 46, rev.	6	
book-licecarpet beetle	5, rev.	4	
cockroaches	51, rev.	12	
greenhouse white fly	57	5-8	
household insects, circular mealy bugs, etc	46, rev. 37, rev. 2	1-7 4	
melon aphis	80	9	
plant-lice	46, rev.	1	
San Jose scale in nursery stock	42, ed. 5 37, rev. 2	7 2	
scale ( Chionaspis sp.)	46, rev.	1	
pests in greenhouses and cold			
frames	37, rev.	1 2	
stored-product insects	46, rev. 46, rev.	1	
white ants	50, rev.	8	
fly	46, rev.	1	
tailed mealy bugwoolly aphis in nursery stock	37, rev. 2	3 6	
fumigation of greenhouses and cold frames nursery stock against San Jose	37, rev. 2	1-11	1-8
Undertiness charges circular	42, ed. 5	7 1–5	1-4
Hylastinus obscurus, circularremedies	67	4-5	
Hylesinus granulatus, injury to lowland fir.	61	9	
n. sp., injury to lowland fir	61	9 2 10	
Western hemlock	61 35	2, 3, 10 $8$	
Hyoscyamus, food plant of Leptinotarsa decemlineata	87	8	
Hymenopterous parasites of house-fly larvæ Hyoscyamus, food plant of Leptinotarsa decemlineata Hyperacmus tineæ, parasite of Tinea pellionella Hyperaspis binotala, enemy of Pulvinaria innumerabilis signata, enemy of Pulvinaria innumerabilis undulata, enemy of Toxoptera graminum Hunhantia eunem	36. rev.	4	
Hyperaspis oinotata, enemy of Pulvingria innumerabilis	64 64	3,5	1
undulata, enemy of Toxontera graminum	93. rev.	6	
		0.0	0.0
Hypoderma bovis, adult, puparium.  lineata, circular.	25 25	2, 8 1–10	3,9
remedies.	25	8-10	
Hypopteromalus tabacum, parasite of Microplitis (Apanteles) catalpæ.	96	5	
I.			
Ichneumon flies, enemies of canker-worms	9	2	
daho, requirements relative to shipment of nursery stock	75, rev. 2	3	
Illinois, requirements relative to shipment of nursery stock	75, rev. 2 59	3	
Index to Circulars 1 to 100 of the Bureau of Entomology.	100	i-ii, 1-49	
Indiana, requirements relative to shipment of nursery stock	75, rev. 2	3	
Insecticide apparatus (see also Mechanical methods).	18	-	
against boll weevil	27	5	
	33	5-6	
cockroaches	51, rev.	13-14	
cockroaches Colorado potato beetle	87	12	

	Circular No.	Page.	Figure.
Insecticide apparatus against elm leaf-beetle	. 8	4	
grasshoppers	. 84	8-9	1
hop plant-louse	. 2	6-7	
melon aphis	80	13-15	5-6
pea aphis	43, ed. 2	7-9	5-6
plum curculio root maggots	63, ed. 2	6–8 5	4-
San Jose scale	3	8	
shade-tree insects	15	2-3	
tobacco thrips	. 68	- 4	
vellow-winged locust	. 53	2-3	
for fumigating greenhouse plants	. 37, rev. 2	9-10	
condensed information concerning some of those most	. 1	6-7	
important, circular	1	1-7	
	16 14	5	
ineffective against boll weevil	18	6	
nsect lime. (See Raupenleim and Dendrolene.)			
nsects defoliating shade trees, control in cities and towns, circular.	- 15	1-4	
dried, food of Tineola biselliella	36	4	
subterranean, insecticides	77	5	
own requirements relative to shipment of nursery stock	75 row 9	6 3	
odine as remedy for "chiggers" owa, requirements relative to shipment of nursery stock socratus vulgaris, parasite of Macrosiphum pisi.	75, rev. 2 43, ed. 2	5	
sosoma tritici, circular	66, rev.	1-7	1-
remedies	66, rev.	. 5–7	
tch mite of fowls. (See Cnemidocoptes mutans.)			The state of the s
vy, ground. (See Nepeta glechoma.) poison, food plant of Pulvinaria innumerabilis			
poison, food plant of Pulvinaria innumerabilis	- 64	2	
(xophorus glaucus, food plant of Toxoptera graminum	. 93, rev.	6	
(Setaria), possible lood plant of Toxoptera graminum	- 85	4	
J			
Towards and italia furnication with hadronomic said one	07 0	0	
Vacaranda ovalifolia, fumigation with hydrocyanic-acid gas	. 37, rev. 2	3	
Jamestown weed. (See Datura.) Janthinosoma posticata	. 72	3	1
sayi, distinctive characters of larva		3	
varipes, distinctive characters of larva	72	3	
farring against boll weevil	18	7	
Colorado potato beetle	. 87	12	
hazelnut weevil	. 99	15	
plum curculio	. 73	6,7-9	1
rose-chafer	- 11	. 4	
spring canker-worm. generally impracticable against chestnut weevils.	9 99	2 9	
'Jelly,' meaning of term as applied to beef and hides	25	1	
'Jigger,' colloquial name for harvest mite	77	1	
Jimson weed. (See Datura.)		1	
TOHNSON, FRED, and GIRAULT, A. A., circular, "The Plum Curculio (Conotrachelus nenuphar Herbst)"			
(Conotrachelus nenuphar Herbst)"	- 73	1-10	1-
Joint-worm. (See Isosoma tritici.)	21 0	0	
Juneberry, food plant of Diabrotica vittata	. 31, rev. 2	2 3	
Saperda candidacretata.	32, rev. 3 32, rev. 3	8	
Scolytus rugulosus	29, rev.	2	
	20,2111		
Κ.			
Kainit against rose-chafer	. 11	4	
seed-corn maggot	. 63, ed. 2	3	
decoction ineffective against rose-chaier	11, rev.	3	
Kakerlach, Dutch name for Blatta orientalis	51, rev.	1~ 10	
Agie as trap crop for cabbage apms	- 80	15-16	
food plant of Dightotica 12-punctata	80	15 3	1
Kale as trap crop for cabbage aphis.  melon aphis.  food plant of Diabrotica 12-punctata.  Murgantia histrionica.  Kansas, requirements relative to shipment of nursery stock.	. 10	2	
Kansas, requirements relative to shipment of nursery stock	75, rev. 2	3	
	. 75, rev. 2	3-4	
Kentucky, requirements relative to shipment of nursery stock	. 47, rev.	ĩ	
Kerosene against bedbug	99	3	
Kerosene against bedbugbox-elder plant-bug.		3	
Xerosene against bedbug.  box-elder plant-bug  carpet beetle.	. 5, rev.		
Kerosene against bedbug.  box-elder plant-bug carpet beetle clover mite	5, rev. 19, ed. 2	4	
Kerosene against bedbug  box-elder plant-bug  carpet beetle.  clover mite  fruit-tree bark-beetle.	5, rev. 19, ed. 2 29, rev.	4 8	
Kerosene against bedbug	5, rev. 19, ed. 2 29, rev. 84	4	
Kerosene against bedbug.  box-elder plant-bug carpet beetle clover mite fruit-tree bark-beetle grasshoppers harlequin cabbage bug.	5, rev. 19, ed. 2 29, rev. 84 10	4 8 8 2	
Kerosene against bedbug.  box-elder plant-bug carpet beetle clover mite. fruit-tree bark-beetle. grasshoppers.	5, rev. 19, ed. 2 29, rev. 84 10	4 8 8	
Kerosene against bedbug.  box-elder plant-bug carpet beetle clover mite fruit-tree bark-beetle grasshoppers harlequin cabbage bug house flies.	5, rev. 19, ed. 2 29, rev. 84 10 { 35, 71, rev. 34, rev.	4 8 8 6,7 6-7 2	
Kerosene against bedobg.  box-elder plant-bug carpet beetle clover mite. fruit-tree bark-beetle. grasshoppers harlequin cabbage bug house flies. little red ant locust borer.	5, rev. 19, ed. 2 29, rev. 84 10 35, 71, rev. 34, rev.	4 8 8 6,7 6-7 2 6	
Kerosene against bedbug.  box-elder plant-bug carpet beetle clover mite fruit-tree bark-beetle grasshoppers harlequin cabbage bug house flies little red ant locust borer mosquitoes in houses	5, rev. 19, ed. 2 29, rev. 84 10 35, 71, rev. 34, rev. 83 13	4 8 8 2 6,7 6-7 2 6 2	
Kerosene against bedbug.  box-elder plant-bug carpet beetle clover mite. fruit-tree bark-beetle grasshoppers harlequin cabbage bug house flies little red ant locust borer.	5, rev. 19, ed. 2 29, rev. 84 10 35, 71, rev. 34, rev. 83 13	4 8 8 6,7 6-7 2 6	

	Circular No.	Page.	Figure.
Kerosene against round-headed apple-tree borer	32, rev. 3	7	
slender seed-corn ground-beetle	78	5	
San Jose scale	42, ed. 5	4-5	
strawberry weevilstriped cucumber beetle	21, rev. 31, rev. 2	. 7	
two-lined chestnut borer	24, rev.	7	
white ants	50, rev.	8	
and carbolic acid against common hen louse	92	6	
gypsum against squash bug. land plaster against squash bug	39, rev. 2 39, rev. 2	5 5	
naphthaline against common hen louse	92	6-7	
plaster ineffective against strawberry weevil	21, rev.	.7	
sand against corn root-aphis	86	12 2	
seed-corn maggot	63, ed. 2 11, rev.	3	
emulsion against apple aphides	81	9,10	
box-elder plant-bug	28	3	
buffalo tree-hopper	23	4	
chicken mite	19, ed. 2	3 4	
cotton red spider	65	5	
cottony maple scale	64	5-6	
elm leaf-beetle	8, rev.	6	
"green bug" harlequin cabbage bug hop plant-louse.	93, rev.	7,19	
hop plant-louse	10	5–6	
imported cabbage worm	60	7	
locust borer	83	5	
melon anhis	80	13-14	
pea aphis pear-tree Psylla rose-chafer	43, ed. 2	7	
pear-tree Psylla	11 row	7 4	
San Jose scale	11, rev.	6	
squash bugs	80	16	
subterranean insects	1	5	
terrapin scale	88	4	
tobacco thrips	24, rev.	3-4, 5	
woolly aphis	20, rev.	4, 5, 6	
and Bordeaux mixture against pear-tree Psylla	7	8	
as insecticide	1	3	
ineffective against periodical cicada	89	4	
only partially successful against strawberry weevil	21, rev.	7	
useless against chestnut weevils	99	9	
Key to known larvæ of the mosquitoes of the United States, circular.	72	1-6	
Ciln-drying against chestnut weevils	99	11	
L.			
abels eaten by Lepisma saccharina	49	1,2	
abena apicalis, parasite of Unrysobolnris jemorala	32, rev. 3	11	
grallator, parasite of Chrysobothris femorata	32, rev. 3	11	
ace-wing flies, enemies of Schizoneura lanigera	20, rev.	6	
fly, golden-eyed. (See Chrysopa oculata.) ady-beetles, enemies of spring grain-aphis (Toxoptera graminum)	93, rev.	3	
adybird, twice-stabbed. (See Chilocorus bivulnerus.)	, , , , , ,		
adybird, twice-stabbed. (See Chilocorus birulnerus.) two-spotted. (See Adalia bipunctata.) ætilia coccidivora, enemy of Pulvinaria innumerabilis	64	4	
ambsquarters. (See Chenopodium,)	04		
ambsquarters. (See Chenopodium.) and plaster and kerosene against squash bug antana, food plant of <i>Aleyrodes vaporariorum</i> .	39, rev. 2	_ 5	
antana, food plant of Aleyrodes vaporariorum	57	/ <sub>5-9</sub>	
asius niger var. americanus, attendant on Aphis maidi-radicis	86	{ 5–9 4	
aws (see also Legislation).	00	, x	
against house flies.	71, rev.	9-10	
relative to importation of American plants, trees, and fruits	44		
into other countries	41	1-4 1-9	
shipment of nursery stock, circulareaf beetle, elm. (See Elm leaf-beetle.)	75, rev. 2	1-9	
rollers of strawberry, arsenicals as remedy	21, rev.	8	
eather, gnawed by cockroaches	51, rev.	4	
ebia atriventris, enemy of Leptinotarsa decemlineatagrandis, enemy of Leptinotarsa decemlineata	87	10	
grandis, enemy of Leptinotarsa decemlineata	87	10	
egislation (see also Laws). against boll weevil	14	7-8	1
emon oil and wood alcohol against corn root-aphis	86	. 13	
Lepidoplatus squamiger	72	6	
episma domestica, frequenter of ovens and fireplaces	49	1, 2, 3	
saccharina, circular	49 49	1-4 3-4	1
			1
entinotarsa decembineata circular	87	1-10	
saccius (ma, circulas, remedies, remedies, eptinotarsa decemlineata, circular, remedies, intermedia	87 87	1-15 12-15	

	Circular No.	Page.	Figure,
Leptocoris trivittatus, circular.	28	1-3	1
remedies. Leptotrachelus dorsalis, enemy of Isosoma tritici.	28 66, rev.	3 4	
Leptus americanus =Trombidium	77 77	$\frac{2,4}{2}$	2
	77	2	1
Lettuce, food plant of Aleyrodes vaporariorum.  Leucania unipuncta, circular.	57 4	2 1–5	1-3
Leucania uniquatica circumata remedies.  Leucopis nigricornis, enemy of Aphis gossypii.  Pulvinaria innumerabilis.	4 80	3-4	
Pulvinaria innumerabilis	64	4	1.0
Lice and mites on poultry, circular poultry, a sure preventive	92 92	1-8 7-8	1-6
poultry, a sure preventive.  "Licked beef," meaning of term Lime against "green bug". peach-tree borer. pear slug. slender seed-corn ground-beetle.	93, rev.	1 19	
peach-tree borer	54 26	4 6	
slender seed-corn ground-beetle.	78	5	
	31, rev. 2 20, rev.	7	
woolly aphis. ineffective against strawberry weevil air-slaked, against striped cucumber beetle.	21, rev. 31, rev. 2	7 7	
impracticable against house flies	ſ 35	6	
and arsenical against peach-tree borer	17	5-6 4	
carbolic acid against common hen louse glue against peach-tree borer	92 17	6 3	
Paris green against two-lined chestnut borer sulphate of copper ineffective against strawberry weevil	24, rev.	7	
sulphur against chicken mite.	21, rev. 92	7 3	
carbolic acid, soap, Paris green, and clay against peach-tree	92	6 4	
borercaterpillar. (See Raupenleim and Dendrolene.)	54	4	
compounds ineffective against rose-chafer	11, rev.	3	
milk, with arsenicals against apple-tree tent caterpillarslaked, against spring grain-aphis	98 85	8 7	
sulphur, and salt wash (see also Lime-sulphur wash).	( 3	7	
against San Jose scale	52, ed. 2	1	
West Indian peach scale	52, ed. 2 52, ed. 2	1-8	
sulphur against cotton red spider wash (see also Lime, sulphur, and salt wash).	65	4-5	
àgainst apple áphides. San Jose scale.	81 42, ed. 5	10 2-4	
worthless against terrapin scale.	88	3	
wash against periodical cicada.  Limnerium fugitivum, parasite of Malacosoma americana	89 98	4 5	
Limnerium fugitivum, parasite of Malacosoma americana. Linden, food plant of Chrysobothris femorata. Pulvinaria innumerabilis.	32, rev. 3 64	9 2	
Linen, starched, eaten by Lepisma saccharina.	49	2	
Liobunum dorsatum, enemy of Leptinotarsa decemlineata Lipeurus variabilis on chicken	87 92	11 7	4
Lixus, resistant to evanid fumes	46, rev. 19, ed. 2	2 2	
Locust, black, food plant of Bryobia pratensis	83	1	
breeding and propagating trees resistant to locust horer	83	7-8	
colloquial name for periodical cicada. differential. (See Melanoplus differentialis.) food plant of Pulvinaria innumerabilis.	30	1	
food plant of Pulvinaria innumerabilis	64	2	
pellucid. (See Camnula pellucida.) Rocky Mountain. (See Melanoplus spretus.)		1	
seventeen-year, colloquial name for periodical cicadatwo-striped. (See Melanoplus bivittatus.)	30	1	
yellow-winged. (See Camnula pellucida.)  Lolium perenne, food plant of Toxoptera graminum	85	4	
London purple against army worm	( 6	3 5	
boll weevil	33	. 2,3	
elm leaf-beetle strawberry weevil. and soap against peach-tree borer	8, rev. 21, rev.	5 6	
and soap against peach-tree boreras insecticide.	17	5-7	
Louisiana, requirements relative to shipment of nursery stock	75, rev. 2	4	
Louse, destructive green pea. (See Macrosiphum pisi.) duck, not same as hen louse.	92	5	
goose, not same as hen louse. hen. (See <i>Menopon pallidum.</i> ) pigeon, not same as hen louse	92	5	
	92	5	

	Circular No.	Page.	Figure.
Lucilia cæsar in houses, habits.	. { 35 71, rev.	2,4,5 2-3	4-5
Lyctus planicollis Lydella doryphoræ. (See Phorocera doryphoræ.)	. 55	2-3	5
Lve-sulphur against cotton red spider	. 65	5	
Lye water against apple aphides.  Lysiphlebus citraphis, parasite of A phis gossypii.  cucurbitaphis, parasite of A phis gossypii.  gossypii, parasite of A phis gossypii.	81 80	9	
cucurbitaphis, parasite of Aphis gossypii	. 80 80	6	
minutus, parasite of A phis gossypii	80	.6	
minutus, parasite of Aphis gossypii tritici, hosts introduction experiments against Toxoptera gra-	. 93, rev.	17	
minum	. 93, rev.	7-14	
parasite of Toxoptera graminum	. \ \ 93, rev.	3-5, 15 <b>-</b> 18	5-10
м.			
Machines. (See Insecticide apparatus.)			
Macrodactylus subspinosus, circular	. 11, rev.	. 1-4 3-4	
Macrosiphum pisi, circular	11, rev. 43, ed. 2	1-10	
Macrosiphum pisi, circular	. 43, ed. 2	6-10	4-7
rudbeckix.  Maggot, cabbage (Pegomya brassicx) (see also Maggot, radish).	. 93, rev.	17	
bisulphid of carbon as insecti-			
cide general account	63, ed. 2	5 3-6	
remedies	. 63, ed. 2	4-6	2-5
imported onion (Pegomya capetorum), bisulphid of carbon as insecticide	. 1	5	
general account	. 63, ed. 2	6-7	
remediesradish ( <i>Pegomya brassicæ</i> ) (see also Maggot, cabbage).	63, ed. 2	7	
bisulphid of carbon as insecticide	. 1	5	
seed-corn. (See Pegomya fusciceps.) Magnolias, food-plants of Macrodactylus subspinosus	. 11, rev.	4	
"Mahogany flat," colloquial name for Cimex lectularius	. 47, rev.	2	
Maine, requirements relative to shipment of nursery stock	75, rev. 2	4 1-8	
Malacosoma americana, circular. remedies.	. 98	6-8	
fragilis, tent caterpillar of the Rocky Mountain region. pluvialis, tent caterpillar of the Pacific Northwest	- 98	2	
spp., hosts of Pimpla inquisitor.	98 97	2 6	
Malaria, carriage by mosquitoes	. 40	1	
Mallows, food plants of Thyridonterux enhemeræformis	. 80 97	5 4	
spp., hosts of Pimpla inquisitor.  Malaria, carriage by mosquitoes.  Mallow, food plant of Aphis gossypii  Mallows, food plants of Thyridopieryx ephemeræformis.  Man, eyelashes and bits of toenails eaten off by cockroaches.	. 51, rev.	6	
host of Dermanyssus gailinæ. harvest mites, or "chiggers" Sarcopsylla pentrans. reported as host of Leptocoris trivittatus.	92	1-3	
Sarcopsylla penetrans	. 77	1	
reported as host of Leptocoris trivittatus seldom bitten by Scutigera forceps.	. 28	3	
Mantis, Carolina. (See Stagmomantis carolina.)	. 10		
Manure (see also Fertilizers and Excrement.) against corn root-aphis.	. 86	11-12	
against corn root-aphis. "green bug".	. 93, rev.	6	
breeding place of Musca domestica	$\left\{\begin{array}{c} 35 \\ 71, \text{ rev.} \end{array}\right.$	5 3	
hen, ineffective against strawberry weevil	. 21, rev.	7	
treatment to destroy house flies	. \ \ \ 71, \text{rev.}	5-8 5-9	
Maple, food plant of Chrysobothris femorata.	. 32, rev. 3	9	
Malacosoma americana. Pulvinaria acericola.	. 98 64	2 2	
innumerabilis Thyridopteryx ephemeræformis	. 64	2	
injury by Ceresa bubalus	97	. 4	
leaf pulvinaria. (See Pulvinaria acericola.) scale, cottony. (See Pulvinaria innumerabilis.) soft. (See Acer saccharinum.)	1		
three species, food plants of <i>Eulecanium nigrofasciatum</i>	. 37, rev. 2	2 3	
the Periodical Cicada"	. 45	1-8	
"Cockroaches" "House Ants (Monomorium pharaonis	. 51, rev.	1-14	1-5
et al.)" pharaons	34, rev.	1-4	1-3
et al.)". "How to Control the San Jose Scale" "Report on the Gypsy Moth and the		17	
Brown-Tail Moth, July, 1904"	. 58	1-12	1-2
Brown-Tail Moth, July, 1904'' "The Bedbug (Cimex lectularius L.)" "The Buffalo Tree-hopper (Ceresa bu-	. 47, rev.	1-8	1-3
balus Fab.)"	. 23	1-4	[1-2]

Lary In C. I. circular ((The Clerer Mite / Drughia material			
MARLATT, C. L., circular, "The Clover Mite (Bryobia pratensis Gar-			
man)" "The Hessian Fly (Cecidomyia destructor	19, ed. 2	1-4	1
Say)"" "The House Centipede (Scutigera forceps	12, rev.	1 -4	
Raf.)"	48	1-4	1-2
"The Imported Elm Leaf-beetle (Galeru- cella luteola Müll.)"	8, rev.	1-6	]
"The Lime, Sulphur, and Salt Wash" "The Ox Warble (Hypoderma lineata	52, ed. 2	1-8	
Villers)''	25	1-10	1-10
"The Peach-tree Borer (Sannina exitiosa   Say)"	17 54	1-4	
"The Pear Slug (Eriocampoides limacina			
Retzius)" "The Pear-tree Psylla (Psylla pyricola	26	1 7	1
Foerst.)". "The Periodical Cicada in 1906 (Tibicen	7	1-8	1-6
septendecim			
L.)'' 1907 ( <i>Tibicen</i>	74	1-5	1-3
septendecim L. race			1
tredecim			
Walsh-Riley)"	89	1-4	1-3
"The Silver Fish (Lepisma saccharina Linn.)"	49	1-4	1-5
"The True Clothes Moths (Tinea pellion- ella et al.)"	36, rev.	1-8	1-0
"The White Ant ( Termes flavipes Koll.)"	50, rev.	1-8	1
"The Woolly Aphis of the Apple (Schizo- neura lanigera Hausmann)"	20, rev.	1-6	1-5
Maryland, requirements relative to shipment of nursery stock	75, rev. 2 75, rev. 2	4	
Mayetiola destructor (see also Cecidomyia destructor).			
circularremedies	70 70	1-16 13-16	1-16
Meadowlarks, enemies of grasshoppers  Mealy bug, (See Dactylopius adonidum.)  "white-tailed." (See Orthezia insignis.)	84	4	
"Meat fly," colloquial name for Calliphora erythrocephala	35 71, rev.	2 2	
Mechanical methods (see also Insecticide apparatus).			
against army wormox warble	25	3, 4	
peach-tree borer	17 54	$\frac{3,4}{3,5}$	1
rose-chafer.	11, rev.	4	
round-headed apple-tree borertwo-lined chestnut borer	32, rev. 3 24, rev.	6 5-6	1
Medicago sativa. (See Alfalfa.) Megarhinus, distinctive characters of genus, adults	40	4	1
hæmorrhoidalis, distinctive characters of adult	40	$\frac{\hat{7}}{2}$	
portoricensis, distinctive characters of adultlarva	40 72	1	
rutila, distinctive characters of adulttable of North American species, adults	40	$\frac{7}{4}$	
Megilla maculata, enemy of A phis gossypii	80	6-7	3, d, e
Leptinotarsa decemlineata	43, ed. 2	10 5	
Melanoconion atratus	85 72	5 3	
melanurus, distinctive characters of larva	72	3	
Melanoplus bivittatus, enemy of alfalfa	84 84	1-6 1-6	2
host of Mermis albicans	62, rev. 84	3,4	3, 4
host of Mermis albicans.  Melittia ceto. (See Melittia satyriniformis.)  cucurbitæ. (See Melittia satyriniformis.)	62, rev.	3	
satyriniformis, circular	38, rev. 2	1-6	1-2
remedies	38, rev. 2		
bug, colloquial name for Diabrotica vittata	31, rev. 2 80		
louse, colloquial name for Aphis gossypii.  Melons (see also Cantaloupe, Cucurbits, Muskmelon, and Water-	00	1	
melon). food plants of Aleyrodes vaporariorum.	57	2	
	80	4	
A phis gossypii Diabrotica 12-punctata	59	3	

	Circular No.	Page.	Figure.
Menopon pallidum, general account	92	4-7	
remedies	92	6-7	
dercurial ointment against ox warble	25	10.12	
Iercurial ointment against ox warble. Merisus destructor, parasite of Mayetiola destructor. Mermis acuminata, probable adult of Mermis albicans.	62, rev.	12-13	
	62, rev.	1-6	
Mesochorus aprilinus, parasite of Apanteles congregatus. fice, hydrocyanic-acid gas as remedy. fichigan, requirements relative to shipment of nursery stock.	62, rev.	5	
Mesochorus aprilinus, parasite of Apanteles congregatus	96		
lice, hydrocyanic-acid gas as remedy	46, rev.		
Migraecon a micridia, hostorial discoss of Pontin many	75, rev. 2 60		
Micrococcus pieridis, bacterial disease of Pontia rapæ Micromus posticus, enemy of Aphis gossypii	80		
Microplitis catalpæ, parasite of Ceratomia catalpæ.	96		
Microplitis catalpæ, parasite of Ceratomia catalpæ. lignonette, food plant of Pontia rapæ. ligrations of Pontia rapæ. ligratory instinct among cockroaches.	60	4	
figrations of Pontia rapæ	60		
ligratory instinct among cockroaches	51, rev.		
filfoil. (See Achillea millefolium.) filkweed. (See Asclepias.)	47, rev.	1	
fillet. (See Panicum miliaceum and Bromus unioloides.)	07	11	
Milyas cinctus, enemy of Leptinotarsa decemlineata	87	11	
fimicry in Sannina exitiosa	{ 54	1	
finnesota, requirements relative to shipment of nursery stock	75, rev. 2	4	
Int, horse. (See Monarda fistulosa.)		•	
liscible oils against San Jose scale	42, ed. 5		
fississippi, requirements relative to shipment of nursery stock	75, rev. 2		
fissouri, requirements relative to shipment of nursery stock	75, rev. 2	5	
fite, chicken. (See Dermanyssus gallinæ.) clover. (See Bryobia pratensis.)			
depluming. (See Cnemidocoptes gallinæ.)			1
	( 35	8	
enemy of house fly	71, rev.	9	
grasshopper. (See Trombidium locustarum.)			
six-spotted, on oranges	19, ed. 2	1	
lites and lice on poultry, circular. enemies of canker-worms.	92	1-8	1-
on poultry, a sure preventive.	92		
Mochlostyrax erraticus, distinctive characters of larva	72	3	
folasses and arsenical against boll weevil	33	4	
Monarda fistulosa, food plant of Anthonomus signatus	21, rev.		
Monarthrum mali, injury to oak	82	2	
Monomorium minutum, general account	34, rev.	$2, \bar{3}$	
remedies	34, rev.	3	
pharaonis, enemy of Cimex lectularius	47, rev.	7	
general account remedies.	34, rev.	1, 2	
Jontana requirements relative to shipment of nursery stock	75, rev.	5	
forning-glory, food plant of Aphis gossypii.  forring-glory, food plant of Aphis gossypii.  forring-glory, food plant of Aphis gossypii.  forring-glory, food plant of Aphis gossypii.	80	5	
forrill, A. W., circular, "The Greenhouse White Fly (Alegrodes			
vaporariorum Westw.)''	57	1-9	
fosquito netting. (See Screen, wire.) fosquitoes as carriers of malaria			1
fosquitoes as carriers of malaria	40	1	
general account	13	1-3	
of North America, how to distinguish the different spe-	10	1-7	1-
cies, circulartables for determination of genera	40	1-7	1-
and species, adults	40	4-7	
United States, key for determination of larvæ, circu-	10		
lar	72	1-6	
remedies	13	1-3	
oth, Angoumois grain (Sitotroga cerealella), hydrocyanic-acid gas	40		
as remedy	46, rev.	2	
brown-tail. (See Euproctis chrysorrhaa.) "Buffalo." (See Anthrenus scrophularia.)			
case-making clothes (See Tinea nellionella.)			
case-making clothes. (See Tinea pellionella.) gypsy. (See Porthetria dispar.) Mediterranean flour (Ephestia kuehniella), hydrocyanic-acid			
Mediterranean flour (Ephestia kuehniella), hydrocyanic-acid			
gas as remedy	46, rev.	2	
southern clothes. (See Timeola biselliella.)			
tuescale (See Trichophaga tapetzetta,)			
southern clothes. (See Tineola biselliella.) tapestry. (See Trichophaga tapetzella.) tussock. (See Tussock moth, white-marked, and Hemero- campa lewostigma.)			
webbing clothes. (See Tineola biselliella.)			
webbing clothes. (See <i>Tineola biselliella</i> .)  toths, carpet. (See Moths, clothes.)  clothes, circular			
clothes, circular	36, rev.	1-8	
hydrocyanic-acid gas as remedy	46, rev.	1, 2	
remedies	36, rev.	5–8	
fur. (See Moths, clothes.) small, prey of Scutigera forceps. fulberry, food plant of Pubinaria innumerabilis	40		
small, prey of Scuttgera forceps.	48	2	
Turberry, rood plant of Puttinaria innumeratius	64 97	4	
	77.6		
Thyridopteryx ephemerx/ormis  Mullein, food plant of Leptinotarsa decemlineata	87	8	

	Circular No.	Page.	Figure.
Murgantia histrionica, remedies	10	2	
Musca domestica (see also Fly, house).	( 25	-	
and other house flies, remedies	35 71, rev.	5-10	
general account	$\left\{\begin{array}{c} 35 \\ 71, \text{rev.} \end{array}\right.$	1-8 1, 3-10	1
insects mistaken therefor	$\left\{\begin{array}{c} 35 \\ 71, \text{rev.} \end{array}\right\}$	1-3 1-6	
Muscina stabulans (see also Cyrtoneura stabulans). in houses	71, rev.	2,3	
Museums, infestation by <i>Tineola biselliella</i> Mushroom beds cultivated by tropical white ants Muskmelon (see also Cantaloupe, Cucurbits, and Melons).	36, rev. 50, rev.	4 4	
food plant of <i>Melitia satyriniformis</i> . Mustard as trap crop for <i>Aphis brassicæ</i> . food plant of <i>Diabrotica 12-punctata</i> .	38, rev. 2 80 59	15–16 3	
Murgantia histrionica	10	1-2	
Pontia rapætrap crop for cabbage aphis	60	4	
rap crop for cappage apms	80 80	15-16 15	
melon aphis	62, rev.	5	
N.			
Naphtha against clothes moths	36, rev.	7	
Naphthaline against clothes moths	36, rev. 92	6 4	
powder-post beetles	55	5	
powder-post beetles. striped cucumber beetle	31, rev. 2	7	
and kerosene against common hen louse	92	6	
preventive of ill effects from exposure to "chiggers" Nasturtium, food plant of <i>Pontia rapæ</i>	77 60	5 4	
nebraska, requirements relative to shipment of nursery stock	75, rev. 2	5	
Nectarine food plant of Conotrachelus nenuphar	73	4, 5	
Scolytus rugulosus.	29, rev.	2 3	
Nectarophora curcurbitæ, the squash aphis destructor= Macrosiphum pisi	43, ed. 2	1	
destructor= Macrosiphum pisi Negundo negundo. (See Box-elder.)			
Nelumbo, food plant of <i>Diabrotica 18-punctata.</i> Nemorxa leucanix, parasite of <i>Leucania unipuncta</i> Neocosmospora vasinlecta (see also <i>Bacillus tracheiphilus</i> ).	59 4	3 5 2	
Spread by <i>Diabrotica vitiata</i> Nepeta glechoma, food plant of <i>Aphis gossypii</i> Netherlands, laws relative to importation of American plants, trees	31, rev. 2 80	5	
and fruits Netting. (See Screen.)	41	4	
Nevada, requirements relative to shipment of nursery stock	75, rev. 2 75, rev. 2	5	
New Hampshire, requirements relative to shipment of nursery stock. Jersey, requirements relative to shipment of nursery stock	75, rev. 2	5 5	
Mexico, requirements relative to shipment of nursery stock	75, rev. 2	5	
York, requirements relative to shipment of nursery stock Zealand, laws relative to importation of American plants, trees.	75, rev. 2	5-6	
and fruits	29, rev.	$\frac{4}{7}$	
Newspaper barriers against fruit-tree bark-beetleround-headed apple-tree borer	32, rev. 3	6	
two-lined chestnut borer	24. rev.	6	
Nezara hilaris, enemy of Leptinotarsa decemlineata	87	11 12	
Nighthawk, enemy of Leptinotarsa decemlineata.  Nightshade, beaked. (See Solanum rostratum.)		. 3	
Nitrate of soda against seed-corn maggot North Carolina, requirements relative to shipment of nursery stock	63, ed. 2	. 6	
Dakota, requirements relative to shipment of nursery stock	75, rev. 2 75, rev. 2	6	
Dakota, requirements relative to shipment of nursery stock.  Nursery stock, fumigation with hydrocyanic-acid gas	46, rev.	1	
requirements relative to shipment, circular	75, rev. 2	1-9 1-15	1-1
Nut weevils, circular Nymphac candidissima, odora, and rosea, fumigation with hydrocy-			
anic-acid gas	37, rev. 2	3	
0.			
Oak, food plant of Chrysobothris femorata	32, rev. 3 98	9 2	
Monarthrum mali	82	2	
Platypus compositus	82	2	
Pulvinaria innumerabilis. Thyridopteryx ephemeræformis.	64 97	2 4	
searlet. (See Quercus coccinea.)	3.		
white, food plant of Agrilus bilineatus	24, rev.	4	
yellow. (See Quercus velutina.)	59	3	
Oats, food plant of Diabrotica 12-punctata  Euthrips nicotianæ.	68	3	
Leucania unipuncta	4	1	
Siphocoryne avenæ.		3 4	
		4	

	Circular No.	Page.	Figure.
Oberea schaumii, injury to Populus monilifera	23	2	
Ochlerotatus abfitchii, distinctive characters of larva	72	4	
æstivalis, distinctive characters of larva	72	6	
atropalpus, distinctive characters of larva	72	4	
aurifera, distinctive characters of larva auroides, distinctive characters of larva	72 72	4 5	
bimaculatus, distinctive characters of larva	72	4	
canadensis, distinctive characters of larva	72	5	
cantator, distinctive characters of larva	72	5	
cinereoborealis	72	4	
<b>c</b> onfirmatus	72	5	
damnosus, distinctive characters of larva	72	5	
dupreei, distinctive characters of larva	72 72	4 5	
fitchii, distinctive characters of larva	72	4	
impiger, distinctive characters of larva inconspicuus, distinctive characters of larva	72	5	
lativittatus	72	6	
lazarensis, distinctive characters of larva	72	6	
mitchellæ, distinctive characters of larva	72	5	
nivitarsis, distinctive characters of larva	72	5	
pretans, distinctive characters of larva	72	6	
pullatus, distinctive characters of larva	72	5	
punctor, distinctive characters of larva	72	4	
serratus	72	4	
sollicitans, distinctive character of larva. subcantans, distinctive characters of larva.	$\begin{bmatrix} 72 \\ 72 \end{bmatrix}$	5 6	
sylvestris, distinctive characters of larva	72	4	
tæniorhynchus.	72	5	
trichurus, distinctive characters of larva	72	4	
triseriatus, distinctive characters of larva	72	5	
trivittatus, distinctive characters of larva	72	5	
varipalpus, distinctive characters of larva	72	5	
varipalpus, distinctive characters of larva Ceneria dispar. (See Porthetria dispar.) Dhio, requirements relative to shipment of nursery stock.	75, rev. 2	6	
Oil (see also Kerosene and Petroleum). residuum, against house fly	71 2077	. 8	
residuum, against house fly	71, rev. 42, ed. 5	4-7	
Oil-water treatment against San Jose scale	42, ed. 5	5-6	
Oils against bedbug	47, rev.	7	
washes and emulsions against San Jose scale Dil-water treatment against San Jose scale Oils against bedbug ox warble and carbolic acid against ox warble alphur against ox warble Disaloma, requirements relative to shipment of nursery stock.	25	8-10	
and carbolic acid against ox warble	25	. 8	
sulphur against ox warble	25	8	
Oklahoma, requirements relative to shipment of nursery stock	75, rev. 2	, 6	
Okra, food plant of Aphis gossypii	80	4	
Diaprotica 12-punctata.	59	3 2	
Diabrotica 12-punctata.  Dive, food plant of Eulecanium nigrofasciatum  Onion, food plant of Pegomya cepetorum	63, ed. 2	6-7	
fusciceps. maggot. (See Maggot, imported onion.) imported. (See Maggot, imported onion.)	63, ed. 2	2	
thrips. (See Thrips tabaci.)	07	0	
Ononis natrix, food plant of Hylastinus obscurus. Ocencyrtus anasæ, parasite of Anasa tristis.	39, rev. 2	3 4	
Orange and other citrus fruits, food plants of A phis gossypii	80	4	
aphis. (See Aphis gossypii.) white fly. (See Aleyrodes citri.) Orchard grass. (See Dactylis glomerata.)	. 00		
scolytus. (See Scolytus rugulosus.)			
	[ 22 ]	1	
	30	1	
trees, injury by periodical cicada	44	1-2	
	74	5	
	89	4	
Oregon, requirements relative to shipment of nursery stock	75, rev. 2	6	
Oriole, Baltimore, enemy of Ceratomia catalpa.	96 98	6	
Malacosoma americana  Ornamental plants, food of Macrodactylus subspinosus	11, rev.	2	
Ortheria insignis on coleus	37, rev. 2	3	
Osage-orange, food plant of Pulvinaria innumerabilis. Pulvinaria. (See Pulvinaria macluræ.)	64	2	
Otionhamahan augusta minta kan far 4 nthanamaciuræ.)			
Otiorhynchus ovatus, mistaken for Anthonomus signatus as straw-	21 2077	3	
berry pest in Rocky Mountain region.  Ovens, infestation by Lepisma domestica.	21, rev. 49	3	
Oxwarble. (See Hypoderma lineata "wormal," colloquial name for Hypoderma lineata	25	1	
P.			
72 - 7	200		
Pachyneuron sp., parasite of A phis gossypii	80	6	
Del consideration and the second and	9	1-3	1-2
Paleacrita vernata, general account		0.0	
Paleacrita vernata, general account	9	2-3	
Paleacrita vernata, general account		2-3	

	Circular No.	Page.	Figure.
Panicum miliaceum, food plant of Diabrotica 12-punctata	59	4	
possible food plant of Toxoptera graminum	85	4	
Paper, wall, injury by Lepisma saccharina	17	2 3	;
wrappings against peach-tree borer	54	3	
Papers, destruction by Termes flavipes	50, rev.	4, 5	
eaten by Lepisma saccharina.	49	1, 2	
Paragordius varius mistaken for Mermis albicans.  Parasites and enemies of brown-tail moth, importation from abroad.	62, rev. 58	5 12	
gypsy moth, importation from abroad	58	9-10	
encouragement against bagworm	97	_ 9	
pine weevil. introduction against "green bug"	90	7.8	6
protection in work against catalpa sphinx	93, rev. 96	7-14 6-7	
Parharmonia picex, work in Sitka spruce	61	9	
Paris green against apple-tree tent caterpillar	98	8	
army worm	97	0.10	
bagworm	6	9-10 5	
boll weevil	33	2,3	
catalpa sphinx	96	6	
Colorado potato beetle	87	13-14	
corn root-worm, southernelm leaf-beetle.	8, rev.	5 5	
fall canker-worm	9	3	
imported cabbage worm	60	6	
pear slug	26 73	6	• • • • • • • • • • • • • • • • • • • •
plum curculioslender seed-corn ground-beetle	78	9 5	
spring canker-worm	9	2	
strawberry weevii	21, rev.	8	
striped cucumber beetle	31, rev. 2	6	
and Bordeaux mixture against strawberry weevil bran mash against imported cabbage worm	21, rev. 60	8,9	
corn meal against tobacco budworm	68	3-4	
lime against two-lined chestnut borer	24, rev.	7	
plaster ineffective against strawberry weevil	21, rev.	7	
salt and horse droppings against grasshoppers	84	9	
yellow - winged locust	* 53	3	
soap against peach-tree borer	17	4	
as insecticide.	1 17	5-7	
carbolic acid, soap, lime, and clay against peach-tree borer	$\left\{\begin{array}{c c} 17\\54\end{array}\right $	4 4	
originally used as insecticide to destroy Colorado potato	01	-	
beetle	87	2	
useless against chestnut weevils. Parthenogenesis in Lysiphlebus tritici.	99	6 16	
Pasimachus elongatus, enemy of Leptinotarsa decemlineata	93, rev. 87	10	
Paste eaten by cockroaches.	51, rev.	4	
Lepisma saccharina	49	2	
Pasturing against clover root-borer	67	5	
Pea aphis. (See Macrosiphum pisi.) food plant of Diabrotica vittata	31, rev. 2	2	
Hylastinus obscurus	67	3	
Macrodactylus subspinosus	11, rev.	2	
Macrosiphum pisi.	43, ed. 2	1-4	
Pegomyā fusciceps louse, destructive green. (See Macrosiphum pisi.)	63, ed. 2	2	
sweet, food plant of Diabrotica 12-punctata	59	3	
Macrosiphum pisi	43, ed. 2	4	
Peach, food plant of Bryobia pratensis.	19, ed. 2	2 9	
Chrysobôthris femorata. Conotrachelus nenuphar.	32, rev. 3 73	4,5	
Diabrotica 12-punctata	59	3	
Diaspis pentagona	52, ed. 2	4	
Eulecanium nigrofasciatum	- 88	2 1	
Leptocoris trivittatus	28 11, rev.	2	
Malacosoma americana	98	2	
Pulvinaria innumerabilis	64	2	
San Jose scale	30 200	$\frac{1}{2}$	
$Scolytus\ rugulosus$ rosette, quarantine regulations of foreign governments against	29, rev.	2	
introduction from the United States	41	2	
soft scale of Europe. (See Eulecanium persicx.)			
tree borer. (See Sannina exitiosa.)			
tree borer. (See Sannina exitiosa.) yellows, quarantine regulations of foreign governments against introduction from the United States	41	2	
Pear, Bartlett, seriously injured by Psylla pyricola.  Buffon, not seriously injured by Psylla pyricola.	7	2	
D . C	7	2	
Bullon, not seriously injured by Psylla pyricola			
Buffon, not seriously injured by Psylla pyricola.  Duchess, seriously injured by Psylla pyricola.  Dwarf Lawrence, seriously injured by Psylla pyricola.	7 1	$\frac{2}{2}$	

	Circular No.	Page.	Figure.
Pear, food plant of A phis mali  Bryobia pratensis.  Chrysobothris femorata.  Conotrachelus nenuphar.  Diabrotica 12-punctata.  Eriocampoides limacina.  Eulecanium nigrofasciatum.  Euproctis chrysorrhea.  Macrodactylus subspinosus.  Malacosoma a mericana.	81	5	
malifolix	81	7	
Bryobia pratensis.	19, ed. 2	2	
Constracted an account of the Constract	32, rev. 3 73	9 5	
Diahrotica 12-minetata	59	3	
Eriocampoides limacina	26	1-7	
Eulecanium nigrofasciatum	88	2	
Euproctis chrysorthæa	58	11	
Macrodactylus subspinosus	11, rev.	2	
Malacosoma a mericana Psylla pyricola. Pulvinaria innumerabilis.	98	2	
Psylla pyricola	7 64	1-2	
San Jose scale	3	1	
San Jose scale Saperda candida	32. rev. 3	3	
Scolytus rugulosus	32, rev. 3 29, rev.	2	
Siphocoryne avenæ	81	3	
Kieffer, not seriously injured by Psylla pyricola	7 7	2	
LeConte, not seriously injured by Psylla pyricola	7	2	
slug. (See Eriocampoides limacina.)		_	
Standard Lawrence, not seriously injured by Psylla pyricola	7	2	
tree Psylla. (See Psylla pyricola.)	00	10.14	
Pecan, 100d plant of Balaninus caryæ	99	12,14	
Pediculoides ventrioseus, onomy of Jacobna tritici	66 POY	4	
Scôlytus rugulosus.  Siphocoryne avenæ.  Kieffer, not seriously injured by Psylla pyricola.  LeConte, not seriously injured by Psylla pyricola.  slug. (See Eriocampoides limacina.)  Standard Lawrence, not seriously injured by Psylla pyricola.  tree Psylla. (See Psylla pyricola.)  Pecan, food plant of Balaninus caryæ.  weevil. (See Balaninus caryæ.)  Pediculoides ventricosus, enemy of Isosoma tritici.  Pegomya brassicæ. (See Maggot, cabbage, and Maggot, radish.)  cepetorum. (See Maggot, imported onion.)  fusciceps, general account  remedies	66, rev.	4	
cenetorum (See Maggot, Cabbage, and Maggot, Tadish.)			İ
fuscicens general account	63, ed. 2	1-3	
remedies	63, ed. 2	2-3	]
remedies	75, rev. 2	6-7	
Pentilia sp., enemy of Tetranychus gloveri	65	3	
Pepper, food plant of Leptinotarsa decemlineata	87	8	
Pennsylvania, requirements relative to shipment of nursery stock. Pentilia sp., enemy of Tetranychus gloveri. Pepper, food plant of Leptinotarsa decemlineata grass, food plant of Aphis gossypii. Peppermint as remedy for "chiggers" Perillus circumcinctus, enemy of Leptinotarsa decemlineata claudus, enemy of Leptinotarsa decemlineata Malacosoma americana.	80	5	
Peppermint as remedy for "chiggers"	77	6	
Perillus circumcinctus, enemy of Leptinotarsa decemlineata	87	11	
claudus, enemy of Leptinotarsa decemlineata	87	11	
Malacosoma americana	98	6	
Periodical cicada. (See Cicada, periodical.)	51, rev.	1 2 7 9	1-
remedies	51, rev.	1,3,7,8 11-14	1-
Periodical cicada. (See Cicada, periodical.)  Periplaneta americana, general account remedies.  australasiæ, general account.	51, rev.	3,8,9	
remedies	51, rev.	11-14	
germanica. (See Blattella germanica.)	02,1011		
germanica. (See Blattella germanica.) orientalis. (See Blatta orientalis.) Persimmon, food plant of Conotrachelus nenuphar			
Persimmon, food plant of Conotrachelus nenuphar	73	4	
Petroleum against locust borer	83	6	
white ants yellow-winged locust	50, rev.	8	
yellow-winged locust	53	2	
crude, against San Jose scale emulsion, crude, against apple aphides soap emulsion against San Jose scale	42, ed. 5	5	
emulsion, crude, against apple applices	42, ed. 5	9,10	
Potune food plant of Lentinotarea decembinests	42, ed. 5 87	8	
Petunia, food plant of Leptinotarsa decemlineata. Phalangium dorsatum. (See Liobunum dorsatum.) PHILLIPS, E. F., circular, "The Brood Diseases of Bees" Philonthus apicalis, enemy of Leptinotarsa decemlineata.	01	0	
PHILLIPS, E. F., circular, "The Brood Diseases of Bees"	. 79	1-5	
Philonthus apicalis, enemy of Leptinotarsa decemlineata	87	11	
ceparum=Pegomya cepetorum	63, ed. 2	6	
Phormia terrænovæ in houses	71, rev.	2-3	
Phormia terrænovæ in houses. Phorocera doryphoræ, enemy of Leptinotarsa decemlineata	87	11	
Phorodon humuli, circular remedies .	2	1-7	1-
remedies	2	4-7	
Phosphorus paste against cockroaches	51, rev.	12	
Phrynium variegatum, fumigation with hydrocyanic-acid gas.  Phyllobænus dislocatus, enemy of Scolytus rugulosus Phyllobænus deslocatus, enemy of Scolytus rugulosus Phyllosunia germanica. (See Blattella germanica.)  Phyllosticta catalpæ, a leaf-spot disease of catalpa.	37, rev. 2 29, rev.	6	
Phyllodromia germanica (Soo Plattella germanica)	23, 161.	0	
Phyllogicia catalog a leaf-spot disease of catalog	96	6	
Phyllogers of grape (Phyllogera nastatrix) insecticides	1	5	
Phymata wolffii, enemy of Pontia range	60	6	
Phylloxera of grape ( <i>Phylloxera vastatrix</i> ), insecticides	87	8	
Phytolacca decandra, food plant of Tetranychus gloveri	65	3	
food plant of Diabrotica 12-punctata.	59	3	
Phytonomus punctatus, fungus enemy	69	2	
the clover-leaf weevil, work in relation to that			
of clover-flower midge (Dasyneura legumi-	00	1.0	
nicola)	69	1-2	
Picea sitchensis. (See Spruce, Sitka.)			
"Pickle brood." (See Brood, pickle.)	6	2	
"Picudo," Mexican name for Anthonomus grandis	14	5	
Tieudo, Mexican name for Annonomous granus	18	5	
Pieris rang. (See Pontia rang.)	, 10	o o	
Pieris rapæ. (See Pontia rapæ.) Pigeon grass. (See Ixophorus glaucus.)			
- Book of the state of the stat	92	9	
mite, not same as poultry mite			

	Circular No.	Page.	Figure.
Pimpla annulipes, parasite of Malacosoma americana	98	5	
Pimpla annulipes, parasite of Malacosoma americana conquisitor, parasite of Malacosoma americana. Thyridopteryx ephemerseformis	98	5	
Thyridopteryx ephemeræformis	97	6	
inquisitor, parasite of Malacosoma americana	98	5	
Thyridopteryx ephemeræformistussock moth and Malacosoma spp	97 97	* 6–7 6	6
pedalis, parasite of Malacosoma americana	98	5	
Pin borer, girdled-cypress. (See Platypus compositus.)			
borers, injury to girdled cypress	. 82	1-3	
Pine destruction by Dendroctonus frontalis	82   24 roy	1-3	
iack host tree of Pissodes strohi	24, rev. 90	3	
lodgepole. (See Pinus murrayana.)	30		
worms, injury to girdled cypress.  Pine, destruction by Dendroctonus frontalis. jack, host tree of Pissodes strobi. lodgepole. (See Pinus murrayana.) pitch, host tree of Pissodes strobi.	90	3	
weevil, white. (See Pissodes strobi.) white, host tree of Pissodes strobi	90	1-7	2-5
yellow. (See Pinus ponderosa.)	0.0	1.4	
Pinhole injury to girdled cypress, circular	82 61	1-4	1
yellow-pine bark maggot	61	9	
murrayana, host tree of bark maggot	61	9	
Dendroctorus sp	61	9	
Dining agdioum operary of Cabinon sung Ignigera	20, rev.	6	
Pissodes strobi, circular	90	1-8	1-6
remedies	90	7-8	
Plantain, food plant of A phis gossypii	80	5	
Plant and the second of the second and secon	51, rev.	2-3	
lice (see also Aphides). hydrocyanic-acid gas as remedylouse, apple-root. (See Apple root-louse and Schizoneura	46, rev.	1	
[amagera]			
products, food of cockroaches	51, rev.	3	
Plaster, land, against striped cucumber beetle	31, rev. 2	6	
products, food of cockroaches Plaster, land, against striped cucumber beetle.	39, rev. 2	5	
impracticable against house flies	35	6	
		6	
of Paris against carpet beetle	5, rev.	3	
cockroaches	51, rev.	14	
and sulphur against cotton red spider	65	4	15
en peresite of Malacosoma americana	70 98	12-13	10
sp., parasite of Malacosoma americana	29, rev.	6	
Platunus compositus, injury to girdled cypress	82	2-3	
Platypus compositus, injury to girdled cypress			1
Plowrightia morbosa, food of Conotrachelus nenuphar	. 73	4	
Plum curculio. (See Curculio, plum.)	10 - 1 0	0	
food plant of Bryobia pratensis	19, ed. 2	2 9	
Chrysobothris femorata	32, rev. 3	4-5	
Diagnis mentagona	52	4	
Diaspis pentagona. Eriocampoides limacina. Eulecanium nigrofasciatum.	26	$1,\hat{3}$	
Eulecanium nigrofasciatum	88	2	
Leptocoris trivittatus	<sup>4</sup> 28	1	
Leptocoris trivittatus	11, rev.	2	
Malacosoma americana	. 98	2	
Phorodon humuli	2	3-4	
Pulvinaria innumerabilis	64	2	
San Jose scale	3	1	
Sannina exitiosa	[ 17 ]	1	
	54	1-2 2	
Scolytus rugulosus	29, rev. 81	3	
Plusia brassicæ. (See Autographa brassicæ.)			
Pneumaculex signifer, distinctive characters of larva	72	3	
Poa pratensis, food plant of Toxoptera graminum	93, rev.	5	
Podabrus rugulosus, enemy of Macrosiphum pisi	43, ed. 2	.5	
Podisus cynicus, enemy of Leptinotarsa decemlineata.	87	11	
maculiventris, enemy of Leptinotarsa decemlineata	87	11	6
modestus, enemy of Malacosoma americana	98	6	
placidus, enemy of Malacosoma americana	98 98	6	
serieventris, enemy of Malacosoma americana.	98	6	
spinosus, enemy of Malacosoma americana	60	6	
=Podisus maculiventris.	87	11	
Poison from centipede bites	48	3	
ivy food plant of Pulvinaria innumerabilis	64	2	
Poisoned bait against grasshoppers	84	9	
Poisoned bait against grasshoppers.  Poisoning from hairs of brown-tail eatarpillars.  Pokeberry. (See Phytolacca decandra.)  Pokeweed. (See Phytolacca.)	58	10	
Pokeweed. (See Phytolacca.)  Polistes pallipes and other wasps, enemies of Pontia rapæ  enemy of Leptinotarsa decemlineata	60	6	

	Circular No.	Page.	Figure.
Polistes rubiginosus, enemy of Leptinotarsa decemlineata		11	
Pollenia rudis in houses, habits	35	2	
Polygmotus hiemalis porosito of Maustiala destructor	( 11, 1ev.	1-2	
Polypogon monspeliensis, food plant of Toxoptera graminum.  Pontia protodice, the southern cabbage butterfly.	70 70	12 11	11
Polypogon monspeliensis, food plant of Toxontera graminum	93, rev.	6	
Pontia protodice, the southern cabbage butterfly	60	3	
		1-8	1-6
remedies	60	6-8	
Poplar (see also Cottonwood).			
Carolina, food plant of Eulecanium nigrofasciatum	88	2	
Bryobia pratensis Malacosoma americana	19, ed. 2	2	
Pulvinaria innumerabilis		$\frac{2}{2}$	
Thyridopteryx ephemeræformis	64 97	4	
Populus monilifera. (See Cottonwood.)	91	4	
Populus monilifera. (See Cottonwood.) Porter's chess. (See <i>Bromus porteri.</i> ) Porthetria dispar, means of control.			
Porthetria dispar, means of control	58	7-10	
report on conditions in New England, July, 1904.	58	1-10	1-2
Porto Rico, requirements relative to shipment of nursery stock	75, rev. 2	7	
Potato beetle, Colorado. (See Leptinotarsa decemlineata.)	1,		
Potato beetle, Colorado. (See Leptinotarsa decemlineata.) bug, colloquial name for Colorado potato beetle	87	1	
food plant of blister beetles	87	3	
food plant of blister beetles.  Diabrotica 12-punctata.	59	3	
Epicauta vittaia Leptinotarsa decemlineata. Pegomya fusciceps.	87	3	
Leptinotarsa decemlineata	87	1-10	
Pegomya jusciceps	63, ed. 2	2	
of Termes flavipes Potentilla canadensis, food plant of Anthonomus signatus	50, rev.	6	
Poultry (goo also Forms and Chiefren)	21, rev.	2	
Poultry (see also Fowls and Chicken). houses, infestation by Cimex lectularius	47 marr	0	
mites and lice, circular		3	1.0
Powder-post beetles, remedies	92 55	1-8	, 1–6
injury to seasoned wood products, circular	55	3–5 1–5	
Praon cerasaphis, parasite of Macrosiphum pisi	43, ed. 2	5	1
'Preussen," colloquial name for Blattella germanica.	51, rev.	2	
Primula, food plant of Aleurodes vaporariorum	57	2	
Privy, box, breeding place of house flies	71, rev.	8	
Privy, box, breeding place of house flies.  Procedular milberti, enemy of Leptinotarsa decemlineata.	87	11	
Promachus saperdx = Cenocalius populator Prune, food plant of Bryobia pratensis.	32, rev. 3	5	
Prune, food plant of Bryobia pratensis	19, ed. 2	2	
Pruning against apple aphides	81	9	
Prunus simonii, food plant of Eulecanium nigrofasciatum	88	2	
Psorophora, distinctive characters of adults	40	4	
synonyms	72	1	
howardi, distinctive characters of larva	40 72	7	
Psylla, pear-tree. (See Psylla pyricola.)	12	1	
pyricola, circular	7	1-8	1-6
remedies	7	7-8	1 0
simulans = winter form of Psylla pyricola. Pteromalus bimaculatus, parasite of Scolytus rugulosus. sp., parasite of Malacosoma americana.	7	3	
Pteromalus bimaculatus, parasite of Scolytus rugulosus	29, rev.	5	
sp., parasite of Malacosoma americana	98	5	
puparum, parasite of Pontia rapæ	60	6,8	5
Publications of the Bureau of Entomology, list, circular	76, rev. to	1-32	
	Feb. 1,		
Pulex irritans, not the common species in houses in the United States.	1910.	0	
serraticeps, general account.	13	3 3–4	
remedies.	13	3-4	
Pulvinaria acericola, the maple-leaf Pulvinaria	64	2	
innumerabilis, circular	64	1-6	1-4
remedies	64	4-6	
maclurx, the osage-orange Pulvinaria	64	2	
Sumpkin (see also Cucurbits).		Į	
food plant of Anasa tristis	39, rev. 2	1	
Diabrotica 12-punctata	59	3	
longicornis	59	7	
vittata	31, rev. 2	2	
Melittia satyriniformis	38, rev. 2	1,2	
Purslane, food plant of Aphis gossypii.	19, ed. 2	5	
cockroaches	51, rev.	1-2_12	
fleas	13 13		
house centipede.	48		
imported cabbage worm	60		
melon aphis	80		
mosquitoes	13		
silver fish	49	4	
silver fishstriped cucumber beetle	31, rev. 2	7	
		3	
ineffective against rose-chafer. strawberry weevil.	11, rev. 21, rev.	7	

	Circular No.	Page.	Figure.
Q.			
Quail, enemy of boll weevil.	27 84	6 4	
Leptinotarsa decemlineata and chinch bug.	87	11-12	
QUAINTANCE, A. L., circular, "The Aphides Affecting the Apple"	81	110	1-8
grasshoppers.  Leptinotarsa decemlineata and chinch bug  QUAINTANCE, A. L., circular, "The Aphides Affecting the Apple" "The Apple-tree Tent Caterpillar (Malacosoma americana Fab.)"	98	1-8	1-4
Quassia decoctions ineffective against rose-chaier	11, rev.	3	
Quedius molochinus, enemy of Leptinotarsa decemlineata	87		
Quercus coccinea, food plant of Agrilus bilineatus velutina, food plant of Agrilus bilineatus	24, rev. 24, rev.	4 4	
Quince, food plant of Diabrotica 12-punctata.  Eulecanium nigrofasciatum.	59	3	
Eulecanium nigrofasciatum San Jose scale	88	2	
Samerda candida	32 rev 3	3	
Scolytus rugulosus Siphocoryne avenæ	29, rev.	2	
Siphocoryne avenæ	. 81	3	
R.			
Radish, food plant of Murgantia histrionica	10		
Pegomya brassica	63, ed. 2	3	
Pegomya brassica:  Pegomya brassica:  fusciceps:  Pontia rapæ:  maggot. (See Maggot, radish, and Maggot, cabbage.)  Ragweed, food plant of A phis maidi-radicis  Diabrotica longicornis.	63, ed. 2 60	2 4	
maggot. (See Maggot, radish, and Maggot, cabbage.)	30	_	
Ragweed, food plant of Aphis maidi-radicis.	. 86 59	6 7	
I RUIT GODIET UL EDRETITET ÆTOT TITOS	97	. 4	
Toxoptera graminum	93, rev.	6	
Rain. (See Weather.)  Rape butterfly colloquial name for Pontia range	60	1	
Rape butterfly, colloquial name for <i>Pontia rapæ</i> .  trap crop for melon aphis	80	15	
Raphitelus maculatus parasite of Scolutus rugulosus	29, rev.	6,7	5
food plant of Diabrotica 12-nunctata.	21, rev. 59	2 3	
Raspberry, black-cap, food plant of Anthonomus signatus.  food plant of Diabrotica 12-punctata.  Rats, hydrocyanic-acid gas as remedy.  Raupenleim against two-lined chestnut borer.	46, rev.	2 7	
Raupeniem against two-lined chestnut borer.  Redbud tree. (See Cercis canadensis.) cultivated. (See Cercis chinensis.)	24, rev.	7	
Red bug, colloquial name for harvest mite or "chigger".	77	1	
coat, colloquial name for Cimex lectularius. spider, confusion with clover mite.	47, rev. 19, ed. 2	2	
cotton. (See Tetranychus gloveri.) spiders on Phytolacca decandra and Xanthium spp.	10,00. 2	1	
spiders on Phytolacca decandra and Xanthium spp	65	3	
"Reptile," colloquial name for Mermis albicans.	50, rev. 62, rev.	8 2	
Redwood, California, not injured by white ants. "Reptile," colloquial name for Mermis albicans. Resin wash against aerial form of woolly aphis	20, rev.	4	
grape Phylloxera and apple root-louse	1 3	5	
washes against San Jose scale as insecticides.  Rhizophagus bipunctatus, enemy of Scolytus rugulosus.	1	3-5	
Rhizophagus bipunctatus, enemy of Scolytus rugulosus	29, rev.	6	
Rhode Island, requirements relative to shipment of nursery stock	75, rev. 2 98	7 5	
Rhogas intermedius, parasite of Malacosoma americana. Rhubarb, food plant of Diabrotica 12-punctata. Rhyncholophus parvus, enemy of Macrosiphum pisi. [Rhey, C.V.], circular, "Condensed Information Concerning Some	, 59	3	
Rhyncholophus parvus, enemy of Macrosiphum pisi	43, ed. 2	5	
of the More Important Insecticides"	1	1-7	
of the More Important Insecticides" "The Hop Plant-louse and the Remedies to be Used Against It"			1.0
Rivoltasia bifurcata on fowls.	92	1-7	1-6
Roach. (See Cockroach.)			
Road dust. (See Dust.) Robin, enemy of Leptinotarsa decemlineata	87	12	
Robinia pseudacacia. (See Locust, black.) Rocky Mountain locust. (See Melanoplus spretus.)	01	12	
Rocky Mountain locust. (See Melanoplus spretus.)	/ 05		
Rolling against "green bug"	85 93, rev.	6 19	
Root borers of strawberry, arsenicals as remedy	21, rev.	8	
maggots and how to control them, circular	63, ed. 2	1-7	1-5
Rose, food plant of Aleyrodes vaporariorum.	57	2	
Diabrotica 12-punctata. Macrodactylus subspinosus.	59	3 1–2	
Malacosoma americana.	11, rev. 98	2	
Pulvinaria innumerabilis	64	2	
San Jose scale.  Roses, fumigation with hydrocyanic-acid gas	37, rev. 2	1 3	
white, attractive to rose-chafer	11, rev.	4	
	80	5	
Runex, food plant of Aphis gossypii.			
Rumex, food plant of Aphis gossypii "Russen," colloquial name for Blattella germanica	51, rev. 65	$\frac{2}{1}$	
Rumex, food plant of Aphis gossypii "Russen," colloquial name for Blattella germanica. "Rust mite," colloquial name for Tetranychus gloveri. Rusts, food of Diabrotica longicornis	51, rev.		

	Circular No.	Page.	Figure.
Rye, food plant of Diabrotica 12-punctata	59	3,4	
Siphocoryne avenæ		3	
Toxoptera graminum	93, rev.	2,5	
slender wild. (See Elymus striatus.)			
S.			
Sabethinæ, distinctive characters of larvæ	72	1	
Saleratus. (See Soda.) Salt against corn root-aphis	86	12	
grass. (See <i>Distichlis spicata</i> .) Paris green, and horse droppings against grasshoppers		9	
water, hot, against nut weevils	53 99	3 10	
water, hot, against nut weevils.  Salvia, food plant of Aleyrodes paporariorum.	77	5	
Salvia, food plant of Aleyrodes vaporariorum	57	2	
Sand and kerosene against seed-corn maggot bur. (See Solanum rostratum.)	63, ed. 2	2	
flea. (See Sarcopsylla penetrans.)			
flea. (See Sarcopsylla penetrans.) SANDERS, J. G., circular, "The Cottony Maple Scale (Pulvinaria innumerabilis Rathvon)"	0.4		
"The Terrapin Scale (Eulecanium nigro-	64	1-6	1-
"The Terrapin Scale (Eulecanium nigro- fasciatum Pergande)"	88	1-4	1-
Sannina exitiosa, circulars	) 17	1-4	
	17	1-6 3-4	
remedies	54	3-6	
opalescens, pest on Pacific coast	54	1 2	
remedies	32, rev. 3 32, rev. 3	1-7 5-7	
cretata, brief account	32. rev. 3	7-8	
remedies	32, rev. 3 84	8	
Sarcophaga cimbicis, parasite of grasshoppers georgina, parasite of grasshoppers. hunteri, parasite of grasshoppers. Sarcopsylla penetrans, the pernicious sand-flea of tropical America.	84	4-5	
hunteri, parasite of grasshoppers	84	4	
Sarcopsylia penetrans, the pernicious sand-nea of tropical America	77	1 10	
Scalding against chestnut weevils.  Scale, cottony maple. (See Pulvinaria innumerabilis.) insects on citrus trees, hydrocyanic-acid gas as remedy pests in greenhouses and cold frames, hydrocyanic-acid gas as	46, rev.	1	
remedy	46, rev. 41	1 4	
red. (See Aonidia aurantii.) San Jose (Aspidiotus perniciosus), an important enemy to fruit trees, circular	3	1-10	1-
quarantine regulations of foreign governments against introduction on			
American plants, trees, and fruits	41	1, 2, 4	
remedies	$\left\{\begin{array}{c}3\\42,\mathrm{ed.}5\end{array}\right.$	6–10	
resin wash as insecticide	1	4	
terrapin. (See Eulecanium nigrofasciatum.) yellow. (See Aonidia aurantii.) 'Scaly-leg'' of fowls, caused by the "itch mite" Cnemidocoptes	* Valuation in the last of the		
mutans	92	3	
Scenopinus fenestralis, in houses, habits	35	2-3 2-3	
Scent glands of cockroaches	71, rev. 51, rev.	2-3 5	
cheele's green against apple-tree tent caterpillar	98	8	
Colorado potato beetle	62, rev.	14	
Schizoneura lanigera (see also Apple root-louse). enemy of apple.		1	
general account. remedies. 'Schwaben,'' colloquial name for Blattella germanica	20, rev.	1-6	1-:
remedies	20, rev.	4-6	
Schware, Colloquiat name for Blattella germanica. Schware, E. A., circular, "The Periodical Cicada in 1897" "The Periodical Cicada in 1898"	51, rev. 22 30	1-4 1-3	
cirpus, food plant of Diabrotica 12-punctata	59	4	
rugulosus, circular	29, rev.	1-8	1-3
remedies	29, rev.	6-8	
creening against house flies	35 71, rev.	5	
mosquitoes	. 13	1	
Screen, wire, against peach-tree borer	17	3	
round-headed apple-tree borer	32, rev. 3	6	
Screw-worms, attracted to wounds of North American fever tick on			
sheep	91	3	

	Circular No.	Page.	Figure.
Scymnus caudalis, enemy of A phis gossypii	80	6	
cervicalis, enemy of A phis gossypii	80 20, rev.	6	
terminatus, enemy of Aphis gossypii.  Seed corn maggot. (See Pegomya fusciceps.)	80	6,7	3, h, i,
treatment before planting, to repel corn root-aphis. Semiotellus chalcidiphagus. (See Homoporus chalcidiphagus.) Senecio, food plant of Diabrotica 12-punctuta.	86	12	
Senecio, food plant of Diabrotica 12-punctata	59 71, rev.	3,6	
"Serpent," colloquial name for Mermis albicans	62, rev.	3,0	
Seneio, 100d plant of Diaprotica 12-punctata.  Sepsis violacca in houses.  "Serpent," colloquial name for Mermis albicans.  "Seventeen-year locust," colloquial name for periodical cicada.  Shade tree insects, control in cities and towns, circular.  trees, food plants of Bryobia pratensis.  injury by buffalo tree-hopper.	30	1 1	
trees food plants of <i>Bruohia pratensis</i>	15 19, ed. 2	1-14	
injury by buffalo tree-hopper.  Shaking treatment against American and European foul brood  Sharpshooter, glassy-winged. (See Homalodisca coagulata.)	23 79	1-3	
	( : 14	5	
incorrect name for Anthonomus grandis	18	5	
Sheep, host of Boophilus annulatus, circular Shepherd's purse, food plaut of Aphis gossypii Ship biscuits, destruction by cockroaches.	91	1-3	
Shepherd's purse, food plant of Aphrs gossypn	51, rev.	5 5	
Snot-dorer, colloquial name for Scottius rugutosus	29, rev.	1	
Shrew, short-tailed. (See <i>Blarina brevicauda</i> .) Shrike, loggerhead, enemy of grasshoppers.  Sigalphus curculionis, parasite of Conotrachelus nenuphar.	84	4	
Sigalphus curculionis, parasite of Conotrachelus nenuphar	73	6	
var. rufus, parasite of Conotrachelus nenuphar. tibiator, parasite of Anthonomus signatus	73 21, rev.	6 4	
Silk, damage by Lepisma saccharina	49	2	
Silk, damage by Lepisma saccharina.  Silver fish. (See Lepisma saccharina.)  Simplocaria niiida, association with Cheilosia alaskensis  Simpson, C. B., circular, "The Yellow-winged Locust (Camnula pellucida Scudd.)".  Sinea diadema, enemy of Leptinotarsa decemlineata  Siphocoryne avenæ, enemy of apple general account remedies	61	8	
pellucida Scudd.)"	53	1-3	]
Sinea diadema, enemy of Leptinotarsa decemlineata	87	11	
siphocoryne avenx, enemy of apple	81 81	3-5	
remedies. Siphonophora citrifolii. (See Aphis citrifolii.)	81	9-10	
Siphonophora citrifolii. (See Aphis citrifolii.) cucurbitæ. (See Nectarophora cucurbitæ.)	0.4		
Sirups, infestation by house ants	26 POV	1 4	
"Skein" centipede, colloquial name for Scutigera forceps. Skins, food of Trichophaga tapetzella Skunk, enemy of grasshoppers.	48	i	
Skins, food of Trichophaga tapetzella	36, rev. 84	5 4	
Ledinotarsa aecemuneata	87	12	
"Slug." colloquial name for larva of Lentinotarsa decemlineata	87	3	
Eriocampoides limacina	26 26	1 1	
Slugs of strawberry, arsenicals as remedy	21, rev.	8	
Slugs of strawberry, arsenicals as remedy.  Smartweed, food plant of A phis maidi-radicis.  Diabrotica longicornis.	86	6	
Snake, colloquial name for Mermis albicans	62, rev.	7 2	
Snake, colloquial name for Mermis albicans. enemy of grasshoppers. worm, colloquial name for Mermis albicans.	84	4	
worm, colloquial name for Mermis albicans.  Snakes, enemies of Leptinotarsa decemlineata.	62, rev. 87	12 12	
Snuffboxes, breeding place of Musca domestica.	4 35	5	
Snuffboxes, breeding place of Musca domestica.  Soap and carbolic acid against fruit-tree bark-beetle	29, rev.	7	
peach-tree borer round-headed apple-tree borer	32, rev. 3	3 6	
caustic potash against round-headed apple-tree borer	32, rev. 3	6	
London purple against peach-tree borer	. 17	4	
Paris green against peach-tree borer	32, rev. 3	4 6	
carbolic acid, Paris green, lime, and clay against peach-tree	17	4	
fish-oil (see also Soap, whale-oil).	54	4	
against fruit-tree bark-beetle	29, rev.	5-6	
hop plant-louseround-headed apple-tree borer	32, rev. 3	5-6	
Solutions against melon aphis	.1 80	13-14	
squash bugs wash against fruit-tree bark-beetle.	29, rev.	16	
pear slug	. 26	6	
treatment against San Jose scale washes against two-lined chestnut borer	42, ed. 5	4 7	
wasnes against two-lined chestnut borer woolly aphis whale-oil (see also Soap, fish-oil),	24, rev. 20, rev.	4,5-6	
against apple aphis	. 81	9,10	
against apple aphis cottony maple scale	64	5	
fruit-tree bark-beetle greenhouse white fly	29, rev. 57	5 9	
round-headed apple-free horer	1 32 rev 3	6	
"green bug" ineffective against strawberry weevil.	93, rev.	7,19	
menective against straw derry weevil	21, rev. 13	7 4	

	Circular No.	Page.	Figure.
Soapy water against itch mite of fowls.	92	4	
preventive of ill effects from exposure to "chiggers"	77	5	
Soda and arsenite of lime against Colorado potato beetle	87	14	
soap against round-headed apple-tree borer	32, rev. 3	6	
remedy for "chiggers"	"	0	
neata	87	. 8	
rostratum, food plant of Leptinotarsa decemlineata  Soldier bug, armed. (See Podisus spinosus.) bordered. (See Stiretrus anchorago.)	87	3	
Solidago. (See Goldenrod.)	20, rev.	4	
Sorbus aucumaria, food plant of Aphis sorbi	81	7	
Soot against root form of woolly aphis  Sorbus aweuparia, food plant of Aphis sorbi  domestica, food plant of Aphis sorbi  torminalis, food plant of Aphis sorbi  Sorghum, food plant of Aphis maidis	81	7	
torminalis, food plant of Aphis sorbi	81	7	
Sorghum, food plant of Aphis maidis.	86	1.2	
Diabrotka longicornis Diatræa saccharalis [larger corn stalk-borer].	59 16	3	
Toxoptera graminum	85	4	
South Carolina, requirements relative to shipment of nursery stock.	75, rev. 2	7	
Dakota, requirements relative to shipment of nursery stock	75, rev. 2	7	
"Spanier," colloquial name for Blattella germanica	51, rev.	2	
"Spanier," colloquial name for Blattella germanica.  Sparrow, chipping, enemy of Malacosoma americana.  English, enemy of Leptinotarsa decemlineata.	98	6	
English, enemy of Leptinotarsa decemtineata	87	12	
Pulvinaria innumerabilis	64   67	3 3	
Smathing mallidge margaite of Chryschothrie femorata	32, rev. 3	11	
simillimus, parasite of Agrilus bilineatus.	24, rev.	5	
Spelt, food plant of Toxoptera graminum	85	4	
Spelt, food plant of Toxoptera graminum  Sphærophoria cylindrica, enemy of Macrosiphum pisi  Sphecius speciosus, range extending northward	43, ed. 2	5	
Sphecius speciosus, range extending northward	97	$\frac{2}{2}$	
sphenophorus, resistant to cyanid rumes	36, rev.	2	
Sphinx, catalpa. (See Ceratomia catalpæ.) catalpæ, same as Ceratomia catalpæ. Spice bush. (See Benzoin benzoin.) Spider plant. (See Cleome.)	96	1	
	87 20, rev.	11 6	
Spike-grass, marsh. (See Distichlis spicata.)  Spiko-grass, marsh. (See Distichlis spicata.)  Spilochalcis mariæ, parasite of Thyridopteryx ephemeræformis.  Spilocryptus extrematus, parasite of Malacosoma americana.  Spinach, food plant of Aphis gossypii.		7	
Spilocraptus entrematus perceito of Malacosoma americana	97 98	5	8
Spingeh food plant of Aphie gossavii	80	5	
	5, rev.	2	
Spiræas, food plants of Macrodactylus subspinosus Spirochæta apis=giant whips of Bacillus larvæ	11, rev.	4	
Spirochæta apis=giant whips of Bacillus larvæ.  Sporotrichum globuliferum, fungous disease of grasshoppers  Spraying apparatus. (See Insecticide apparatus.)  Spruce, destruction by Dendroctonus frontalis	94 84	3-4 5-7	7
Spring apparatus. (See Insecticine apparatus.)	24, rev.	1	3
host tree of Pissodes strobi	90	3	3
Sitka, host tree of Parharmonia piceæ	61	9	
Squash (see also Cymling and Cucurbits). aphis. (See Necturophora cucurbits.)	61	9,10	
bug. (See Anasa tristis.) black. (See Anasa tristis.) food plant of Anasa tristis. Diabrotica 12-punctata			
1000 plant of Anasa tristis	39, rev. 2 59	2 2	
longicornis	59	2,3	
vittata		i	
vittata	31, rev. 2 38, rev. 2	$1,\bar{2}$	
Hubbard, food plant of Diabrotica vittata	31, rev. 2	2	
marrow, food plant of Diabrotica vittatavine borer. (See Melittia satyriniformis.) Stable fly. (See Cyrtoneura stabulans, Muscina stabulans, and Sto-	31, rev. 2	2	
moxys calcitrans.)	07	0	
Stagmomantis carolina, range extending northward Stalk-borer, larger corn. (See Diatrxa saccharalis.) Starch, eaten by Lepisma saccharina	97 49	1,2	
Steam against carpet beetle	5, rev.	4	
powder-post beetles	55	4	
Stegomyia calopus, distinctive characters of larva	72	5	
Stenanthium lindenii, fumigation with hydrocyanic-acid gas	37, rev. 2	3	
Stenomesius aphidicola, parasite of Aphis gossypii	80	6	
Sticky fly paper against house flies	$\begin{cases} 35 \\ 71, \text{ rev.} \end{cases}$	5 5	
Stictonotus isosomatis, parasite of Isosoma tritici	66, rev.	4	
Stink bug, common name for Anasa tristis Stipa, green. (See Stipa viridula.)	39, rev. 2	1	
viridula, food plant of Toxoptera graminum	93, rev.	6	
Stiretrus anchorago, enemy of Leptinotarsa decemlineata	87 35	11	
Stomorys calcitrans in houses, mistaken for Musca domestica	71, rev.	$1, \frac{1}{2}$	2
Stored foods, drugs, and vegetables, eaten by Lepisma saccharina		1,2	

•	Circular No.	Page.	Figure.
Straw barriers against fruit-tree bark-beetle	29, rev.	7	
peach-tree borer	$\left\{\begin{array}{c} 17 \\ 54 \end{array}\right]$	3	
Strawberry crown-girdler. (See Otiorhynchus ovatus.)	( 94	3	
food plants of Aleyrodes packardi	57	4	
A nthonomus signatus A phis gossypii	21, rev. 80	1-5 5	
slugs, arsenicals as remedy.	21, rev.	8	
slugs, arsenicals as remedyvarieties preferred as food plants by <i>Anthonomus signa</i> -			
weevil. (See Anthonomus signatus.)	21, rev.	1,2,6	
white fly. (See Aleyrodes packardi.) wild, food plant of Anthonomus signatus. "Striped bug," colloquial name for Diabrotica vittata Stylopyga orientalis. (See Blatta orientalis.)	21, rev. 31, rev. 2	$\frac{2}{1}$	
Submersion in water against pin borers in cypress and gilms	82	4	
Sugar and borax against ants. Sugar cane borer. (See Diatræa saccharalis.) food plant of Diatræa saccharalis	34, rev.	4	
food plant of Diatræa saccharalis	16	1,3	
Sugars, infestation by house ants	34, rev. 21, rev.	1 7	
potash against seed-corn maggot	63, ed. 2	3	
against cockroaches	51, rev.	12	
cotton red spider. "green bug". and flour against cotton red spider.	65	4 19	
and flour against cotton red spider	93, rev. 65	19	
lime against chicken mite	92	3	
common hen louse	92	6	
oils against ox warbleplaster of Paris against cotton red spider	25 65	8 4	
road dust against cotton red spider	65	4	
"chiggers"	77	5	
"chiggers" dioxid against carpet beetle. fumigation against bedbug.	5, rev. 47, rev.	4 7	
ointment against itch mite of fowls	92	4	
preventive of ill effects from exposure to "chiggers"	77	5	
Sumac, food plant of Macrodactylus subspinosus	11, rev.	4	
Sunflower, food plant of Diabrotica 12-punctatalongicornis	59 59	3 7	
vittata	31, rev. 2	2	
wild, food plant of Diabrotica 12-punctata	59	3	
Sunlight against chicken mite.	36, rev.	3 6	
clothes moths.  Swallows, cliff or eaves, host of Cimex sp.	47, rev.	3	
Switzerland, laws relative to importation of American plants, trees,	34, rev.	1	
and fruits	32, rev. 3	4 9	
Eulecanium nigrofasciatum	88	2	
Pulvinaria innumerabilis	64	_ 2	
Syrphoctonus maculifrons, parasite of Cheilosia alaskensis.	61	7-8	
Syrphus americanus, enemy of Aphis gossypii.  Macrosiphum pisi	80	6	
Macrosiphum pisi	43, ed. 2	5	
fly larvæ, enemies of aphides	81 80	8	
т.	80	0	
Tachina flies, enemies of canker-worms	9	2	
Txniorhynchus perturbans, distinctive characters of larva	72	3	
Tachina flies, enemies of canker-worms fly, red-tailed. (See Nemorxa leucanix.) Txniorhynchus perturbans, distinctive characters of larva. Tanager, scarlet, enemy of Leptinotarsa decemlineata. Tapestry moth. (See Trickophaga tapetzella.) Tar against white ants.	87	12	••••
Tar against white ants	50, rev.	7	• • • • • • • • • • • • • • • • • • • •
paper against carpet beetle clothes moths.	5, rev. 36, rev.	3 6	***************************************
iruit-tree bark-beetle	29, rev.	7	
two-lined chestnut borer	24, rev.	6	
cards against cabbage maggot.  Tares, food plants of Macrosiphum pisi.  Tea, New Jersey. (See Ceanothus.)	63, ed. 2 43, ed. 2	5-6 4	3–5
Telephorus clisiocampæ, parasite of Malacosoma americana	98 67	5 4	
extremes as limiting distribution of Leptinotarsa de-	87	6-7	
cemlineatain relation to Toxoptera graminum	93, rev.	6-7 3-5	
Tennessee, requirements relative to shipment of nursery stock	75, rev. 2	7	
m - 4 - 4			
Tent caterpillar, apple-tree. (See Malacosoma americana.) caterpillars. (See Malacosoma spp.) Tenthredo cerasi, distinct from Eriocampoides limacina			

	Circular No.	Page.	Figure.
Tenthredo limacina = Eriocampoides limacina	26	2	
Termes flavipes. (See Ant, white.)	50, rev.	4	
lucifugus in America	87	11	
Tetramorium cæspitum, feeding on honeydew from A phis gossypii	80	1	
general accountremedies	34, rev. 34, rev.	3-4	3
Tetranychus gloveri, circular	65	1-5	1-2
remedies	65	4-5	
Tetrastichus microgastri, secondary parasite of Pontia rapæ	29, rev.	5 6	
spp., parasites of Dasyneura leguminicola	69	6	:
ranthomelana parasite of Galericella Luteola	8, rev.	4	
Texas, requirements relative to shipment of nursery stock.  Thanasimus sp., enemy of Scolytus rugulosus.  Theobaldia incidens, distinctive characters of larva.	75, rev. 2 29, rev.	7-8	
Theobaldia incidens, distinctive characters of larva	72	4	
Thermobia domestica. (See Lepisma domestica.)			
Theronia melanocephala, parasite of Malacosoma americana	98	6	
Theodadia incidents, distinctive characters of latva. Thermobia domestica. (See Lepisma domestica.) furnorum. (See Lepisma domestica.) Theronia melanocephala, parasite of Malacosoma americana. Therilochus conotracheli, parasite of Conotrachelus nenuphar. Thistol of facotic access	73	6	
I histie, howers nequented by butterines of I onthe rapa	60	4	
food plant of Diabrotica longicornis	59 87	7 8	
Thorn, food plant of Malacosoma americana.	98	2	
Superau cretatu	32, rev. 3	8	
white. (See Cratagus oxyacantha.)	32, rev. 3	3	
Thorns, food plants of Saperda candida "Thousand legs," colloquial name for Scutigera forceps	48		
Thrips, hydrocyanic-acid gas as remedy in greenhouses, remedies.	46, rev.	1	
in greenhouses, remedies	80	12	
tabaci, the onion thrips, not found on tobacco in United			1
Statestobacco. (See Euthrips nicotianæ.)	68	1	
wheat. (See Euthrips tritici.) Thrush, hermit, enemy of Leptinotarsa decemlineata	07	19	-
olive-backed, enemy of Leptinotarsa decembratea	87 87	12	
wood, enemy of Leptinotarsa decemlineata	87	12	
Tibicen septendecim. (See Cicada, periodical.)			
Tick, chicken. (See Argas miniatus.)	77	1	
erroneous name for harvest mite or "chigger"	77	1	
North American fever. (See Boophilus annulatus.) Timber beetles, injury to girdled cypress. Timothy, food plant of Leucania unipuncta.	82	1	
Timothy, food plant of Leucania unipuncta.	70	1	
Mayetiola destructor or related species  Tinea pellionella, general account.	36, rev.	7 2-4	ii
remedies	36, rev.	5-8	
Tineola biselliella, general account	36. rev.	4-5 5-8	2
remedies. Trus, E. S. G., circular "The Cotton Red Spider ( <i>Tetranychus gloveri</i> Bks.)".	36, rev.	5-0	
gloveri Bks.)"	65	1-5	1-2
Toad, enemy of grasshoppers.  Leptinotarsa decemlineata.	84 87	4 12	
Malacosoma americana	98	6	
Tobacco against aphides, etc	37, rev. 2		
apple aphides.	81 36, rev.	10	
common hen louse	92	6	
"green bug". greenhouse white fly	93, rev.	6	
greenhouse white fly	57 80	8,9 10-13	
greenhouse white fly melon aphis. striped cucumber beetle woolly aphis. and soapsuds against "green bug" budworms. (See Chloridea virescens and Heliothis obsoleta.) food plant of Diobratin 12-avanta.	31, rev. 2	7	
woolly aphis	20, rev.	4,5	
and soapsuds against "green bug".	93, rev.	6	
food plant of Diabrotica 12-punctata	59	3	
Euthrips nicotianx	68	1-2	
Leptinotarsa decemlineataineffective against rose-chafer	87 11, rev.	8 3	
strawberry weevil	21, rev.	7	1
thrips. (See Euthrips nicotianx.)	68	1-2	
"white veins" caused by Euthrips nicotians  Tomato, food plant of Aleyrodes vaporariorum	\$ 37, rev. 2	4	
A phis gossypii	57 80	1.2	
Diabrotica 12-punctata.	59	3	
Diabrotica 1?-punctata	87	8	
Tomatoes, fumigation with hydrocyanic-acid gas	37, rev. 2 85	4 1-7	1-3
Toxoptera graminum, circulars	93, rev.	1-22	1-10
*	93, rev.	6-7	
experiments with remedies	85	6-7	,

	Circular No.	Page.	Figure.
Trapping against boll weevil	56	3-4	
cockroaches	51, rev.	13-14	
squash bug	39, rev.	. 4	
Trap crops against buffalo tree-hopper harlequin cabbage bug.	23	4 2	
Hessian fly	12, rev.	4	
imported cabbage worm	60	8	
melon aphis	80	15	
rose-chaferstrawberry weevil	11, rev. 21, rev.	4 5	
useless against chestnut weevils	99	9	
leaves against imported cabbage worm	60	8	
plants against boll weevil	14 18	6,8 6–7	
	95, rev.	5-6	
striped cucumber beetletrees against two-lined chestnut borer	31, rev. 2	5-6	
trees against two-lined chestnut borer	24, rev.	6	
felled, against ambrosia beetles in cypress	32. rev. 3	3-4 11	
Praps against fruit-tree bark-beetle	29, rev.	7	
squash-vine borer	38, rev. 2	5	
Pree-hopper, buffalo. (See Ceresa bubalus.) Prees, living, eaten by Termes flavipes	50 man	F 0	
Trichasis rufipes, probable parasite of Balaninus proboscideus and B.	50, rev.	5–6	
rectus	99	6	
Trichonaga tanetzella, general account	36, rev.	5	3
remedies.  Trichopoda pennipes, parasite of Anasa tristis.	36, rev.	5-8	
Trifolium hybridum. (See Clover, alsike.)	39, rev. 2	4	
pratense. (See Clover, red.) Trioxys testaceipes, parasite of Aphis gossypii. Trioxacur dactyloides, food plant of Diatræa saccharalis (corn stalk-	80	6	
horer)	16	3	
Trissolcus murgantiæ, parasite of Murgantia histrionica Trombidium, Leptus a synonym locustarum, enemy of grasshoppers	10 77	2 2	
locustarum, enemy of grasshoppers	84	4	
sp	77	1,3	1,3
Fubers, starchy, food of cockroaches.	51, rev.	3	
w. t. a.	32, rev. 3 27	10 5	
fruits Furnip, food plant of Diabrotica 12-punctata	41 59	4 3	
Pegomya fusciceps	63, ed. 2	2	
Pontia rapæ	60	4	
Furpentine against bedbug.  ox warble	47, rev. 25	7 9	
striped cucumber beetle	31, rev. 2	7	
and gypsum against squash bug	39, rev. 2	5	
and gypsum against squash bug land plaster against squash bug	59, rev. 2	5	
Fussock moth, white-marked (Hemerocampa leucostigma), control host of Pimpla inquisitor	15 97	3, 4	
shade-tree pest.	97	1	
Tweezers for removal of ox warble	25	9	
Typhoid fever, carriage by house fly	35	8	
fly, proposed as a better name for Musca domestica	71, rev. 71, rev.	8 [11]	
U.	71, 164.	[11]	
Ulmus americana notably exempt from attack of Galerucella luteola	8, rev.	3	
Uranotænia, distinctive characters of larvæ	72	3	
sapphirinaUropoda americana, enemy of Leptinotarsa decemlineata	72 87	3 11	
Urosigalphus armatus, parasite of Balaninus proboscideus and B.	01	11	
rectus	99	6	
Utah, requirements relative to shipment of nursery stock	75, rev.	8	
$\nabla.$			
Vaccinium sp., food plant of Eulecanium nigrofasciatum	88	2	
"chiggers"	77	5	
preventive of ill effects from exposure to "chiggers"	77	5 8	
Vermont, requirements relative to shipment of nursery stock Vetches, food plants of Macrosiphum pisi	75, rev. 2 43, ed. 2	8	
Violet, food plant of Aphis gossypii.	80	5	
Violets, fumigation with hydrocyanic-acid gas. Vireo, red-eyed, enemy of Malacosoma americana	37, rev. 2	3	
Vireo, red-eyed, enemy of Malacosoma americana.  Virginia creeper, food plant of Pulvinaria innumerabilis	98 64	6 2	
TARAMAN CAUCHOL, IOUG PIGHO OL & WOULDER OF FIRE BUILD BUILD	75, rev. 2	8	

	Circular No.	Page.	Figure.
w.			
"Wall louse," colloquial name for Cimex lectularius paper, damage by cockroaches	47, rev.	2	
paper, damage by cockroaches.	51, rev. 25	6	
"Warbled stock," meaning of term Warbler, yellow, enemy of Malacosoma americana	98	6	
Washington [State], requirements relative to shipment of nursery			
stock	75, rev. 2	8	
mimicked by Sannina exitiosa.	§ 17	1	
Wasps, enemies of canker-worms.	54	$\frac{1}{2}$	
feeding on honeydew from aphides	80	1	
Water against cotton red spidermelon aphis	65	5	
holling against elm leal-beetle	s. rev.	14 6	
bug, colloquial name for Biattella germanica	1 51, rev.	1	
		4 7	
bedbug box-elder plant-bug carpet beetle chestnut weevils	47, rev. 28	3	
carpet beetle	5, rev.	3	
chestnut weevils	99 60	10 7	
peach-tree borer	. 54	5	
woolly aphis	20, rev.	4-5, 6	
ineffective against rose-chafer  preventive of ill effects from exposure to "chiggers"	11, rev.	3 5	
preventive of ill effects from exposure to "chiggers" oak, probable food plant of <i>Tetranychus gloveri</i>	65	. 3	
submersion against pin borers in cypress	82	9	
test against chestnut weevils unreliable	99	. 9	
food plant of Diabrotica 12-punctata	. 59	3	
vittata	31, rev. 2	- 2	
Weather (see also Temperature.) in relation to Mayetiola destructor	. 70	10	
Toxoptera graminum	. 85	5-6	
Webster, F. M., circular, "Some Insects Affecting the Production of Red Clover Seed"	. 69	1-9	1-8
"The Clover Root-borer (Hylastinus ob-			10
scurus Marsham)''	. 67	1-5	1-4
"The Corn Leaf-aphis and Corn Root- aphis"	. 86	1-13	1-4
"The Grasshopper Problem and Alfalfa			
"The Hessian Fly (Manetiala [Cecido-	. 84	1-10	1-8
"The Hessian Fly (Mayetiola [Cecido- myia] destructor Say)"	. 70	1-16	1-16
"The Joint-worm (Isosoma tritici Fitch)"	66, rev.	1-7	1-6
"The Slender Seed-corn Ground-beetle	. 78	1-6	1-2
(Clivina impressifrons Lec.)". "The Spring Grain-aphis (Toxoptera graminum Rond.)".			
graminum Rond.)"	. 85	1-7	1-3
"The Spring Grain-aphis or so-called 'Green Bug' (Toxoptera graminum Rond.)".			
Rond.)".	93, rev.	1-22	1-10
Websterellus tritici, parasite of Isosoma tritici. Webworm, fall (Hyphantria cunea), control.	66, rev.	3,4	
on catalpa	. 96	1	
shade-tree pest	97	1 15	
Weeds, eradication against Colorado potato beetle	27	5	
clover-leaf. (See Phytonomus punctatus.)			
hazelnut. (See Balaninus obtusus.) larger chestnut. (See Balaninus proboscideus.)			
lesser chestnut. (See Balaninus rectus.)			
Marrican action hall (Anthonomy arandia) general airculant	6 14	1-5	1-3
$\textbf{Mexican cotton-boll}  (\textbf{\textit{Anthonomus grandis}}), \textbf{general circulars}.$	18	1-8 1-8	1-5
in 1897, circular.	. 27	1-7	
most important remedy, circu-	56	1-7	
lars	.])	1-8	
remedies, circular	33	1-6	
pecan. (See Balaninus caryæ.) white pine. (See Pissodes strobi.)			
West Indian peach scale. (See Diaspis pentagona)	. 52, ed. 2	2	
Virginia, requirements relative to shipment of nursery stock. Wharves, destruction by white ants.	75, rev. 2 50, rev.	8	
Wheat, durum or macaroni, not attractive to Hessian fly	. 70	10	
food plant of (Cecidomyia) Mayetiola destructor	12, rev.	1,2-3	
	70	1-10	
chinch bug. Diabrotica 12-punctata Isosoma tritici	. 59	3, 4 1-4	

	Circular No	Page.	Figure.
Wheat, food plant of Macrodactylus subspinosus		2 4	
Siphocoryne avenæ	81 85	4	
Toxoptera graminum	93, rev.	2,5	
grass, slender. (See Agropyron tenerum.) midge. (See Contarinia tritici.) thrips. (See Euthrips tritici.)			
varieties resistant to Hessian fly	12, rev.	4	
varieties resistant to Hessian fly volunteer, destruction in control of Hessian fly	70	15	
Wheel bug. (See Arilus cristatus.)			
White fly, greenhouse. (See Aleyrodes vaporariorum.)	40		
hydrocyanic-acid gas as remedy orange. (See Alegrodes citri.)	46, rev.	1	
strawberry. (See Aleurodes packardi.)			
White, G. F., circular, "The Cause of American Foul Brood"	94	1-4	
"White veins" in wrapper tobacco the work of Euthrips nicotianx.	68	1-2	
Whitewash and carbolic acid against chicken mite	92	3	
Willow, food plant of Chrysobothris femorata	32, rev. 3	9 2	
Pulvinaria innumerabilis	98 64	2	
injury by Ceresa bubalus	23	2	
Wilt disease of cucurbits, cotton, and cowpeas, bacterial. (See Bacillus tracheiphilus and Neocosmospora vasinfecta. Window fly. (See Scenopinus fenestralis.)			
Wire gauze (see also Screen, wire).	00 0		
barriers against round-headed apple-tree borer Wisconsin, requirements relative to shipment of nursery stock	32, rev. 3 75, rev. 2		
Witch-hazel, food plant of Malacosoma americana.	98	0-9	
Woodpecker, red-headed, enemy of grasshoppers	84	4	
Woodpeckers, enemies of Chrysobothris femorata	32, rev. 3	11	
Pissodes strobi	90	- 7	
Saperda candida	32, rev. 3	5	
Wood, seasoned, injury by powder-post beetles, circular.  Woods, Albert F., circular, "The Use of Hydrocyanic-acid Gas	55	1-5	
for Fumigating Greenhouses and Cold			
Frames"	37, rev. 2	1-11	1-3
and Dorsett, P. H., circular, "The Use of Hydrocyanic-acid Gas for Fumigating			
Hydrocyanic-acid Gas for Fumigating		1 1 2 22	
Greenhouses and Cold Frames"	37, rev.	1-10	1-3
Woolens, food of Anthrenus scrophularix  Tinea pellionella	5, rev. 36, rev.		
Tineola biselliella	36, rev.	4	
Trichophaga tapetzella	36, rev.	5	
gnawed by cockroaches.	51, rev.	4	
Woolly aphis. (See Schizoneura lanigera.)			
apple louse. (See Schizoneura lanigera.) "Worms," colloquial name for chestnut-weevil larvæ	00	2	
Wyeomyia smithii, distinctive characters of larva	99 72	1	
Wyoming, requirements relative to shipment of nursery stock		9	
X.		•	
Vanthium ann (Coo Cooldohum)			
Xanthium spp. (See Cocklebur.) Xanthosoma lindeni, fumigation with hydrocyanic-acid gas	37, rev. 2	3	
zzammoooma macmi, tumigation with nyurocyame-acid gas	01, 10V. Z	. 0	

			The second secon
	7		
			contract and make a second and a second as
			to a construction of the large for the plantage of the contract of the contrac
			The state of the s
			the second secon
			percentification (1984) and the second second
Appropriate to the second seco			
Appropriate to the second seco			
2.70 (A), see the state of			



